arXiv:1110.1520v2 [math.AT] 28 Dec 2011

ARRANGEMENTS OF SUBMANIFOLDS AND THE TANGENT BUNDLE COMPLEMENT

PRIYAVRAT DESHPANDE

Abstract. Drawing parallels with the theory of hyperplane arrangements, we develop the theory of arrangements of submanifolds. Given a smooth, finite dimensional, real manifold X we consider a finite collection $\mathcal{A}$ of locally flat codimension 1 submanifolds that intersect like hyperplanes. To such a collection we associate two posets: the poset of faces and the poset of intersections. We also associate a topological space to the arrangement. The complement of union of tangent bundles of these submanifolds inside the tangent bundle of the ambient manifold which we call the tangent bundle complement. Our aim is to investigate the relationship between combinatorics of the posets and topology of the complement.

Using the Nerve lemma we show that the tangent bundle complement has the homotopy type of a finite simplicial complex. We generalize the classical theorem of Salvetti [21, Theorem 1] for hyperplane arrangements and show that this particular simplicial complex, called the Salvetti complex, is completely determined by the face poset. We also characterize all the connected covers of this complex, thus generalizing the work of Delucchi [7] and Paris [19].

Introduction

An arrangement of hyperplanes is a finite set $\mathcal{A}$ consisting of linear (or affine) codimension 1 subspaces of $\mathbb{R}^l$. These hyperplanes and their intersections induce a stratification of the ambient space. The strata form a poset when ordered by inclusion and the set of all possible intersections forms a poset ordered by reverse inclusion. These posets contain important combinatorial information about the arrangement. The topological objects associated with an arrangement $\mathcal{A}$ are the *real complement* $\mathcal{C}(\mathcal{A})$ and the *complexified complement* $M(\mathcal{A})$. The real complement is the complement of the union of hyperplanes in $\mathbb{R}^l$, whereas the complexified complement is the complement of the union of the complexified hyperplanes in $\mathbb{C}^l$. One of the important aspects of the theory of arrangements is to understand the interaction between the combinatorial data of an arrangement and the topology of these complements. For example, one would like to comprehend to what extent the combinatorial data of an arrangement control the topological invariants, such as (co)homology or homotopy groups etc., of these complements.

In this paper we introduce a generalization of real hyperplane arrangements which we call as the *arrangements of submanifolds of codimension* 1. We consider situations in which finitely many submanifolds of a given manifold intersect in a way that the local information is same as that of a hyperplane arrangement but the global picture is quite different. Intuitively speaking it means that for every point on that manifold there exists a coordinate neighborhood homeomorphic to an arrangement of real hyperplanes. We also introduce an analogue of the complexified complement in this new setting and call it *the tangent bundle complement.*

2010 *Mathematics Subject Classification.* 52C35, 57N80, 05E45.

Key words and phrases. Hyperplane arrangement, Salvetti complex, Nerve lemma, CW-complex .

All the results proved in this paper attempt to answer the following question: *how does the combinatorics of the intersections of these submanifolds help determine the topology of the tangent bundle complement?*

The topological study of the complement of a complexified hyperplane arrangement can be traced back to the work of Arnold on braid groups and configuration spaces. His results led Brieskorn to conjecture that the (complexified) complement of a real reflection arrangement is a $K(\pi, 1)$ space (i.e. its universal cover is contractible). In 1973, Deligne settled this conjecture in [6]. His first step was to parametrize the universal cover of the complement by sequences of adjacent chambers (the galleries in the Coxeter complex). Using the earlier work of Garside [11] on the conjugacy problem for braid groups he showed that these positive paths can be expressed in a particular normal form. This normal form was the main ingredient in proving that the constructed universal cover was contractible. We should also note that it is still an open question whether having a $K(\pi, 1)$ complement is a combinatorial property. Deligne's work has greatly influenced research in hyperplane arrangements and geometric group theory. For example, existence of a bi-automatic structure in finite type Artin groups as proved in [3] and the discovery of Garside groups in [5] are consequences of Deligne's work.

The combinatorial nature of Deligne's arguments led to the search for *combinatorial models for the complement*, i.e., cell complexes built using the combinatorial data of an arrangement that are homotopy equivalent to the complement. One such model, the *Salvetti complex*, was introduced by Salvetti in [21]. He showed that the incidence relations in the underlying face poset are sufficient to build this CW complex. His construction was generalized to complex arrangements in [1] where the authors also showed that the face poset determines the homeomorphism type of the complement. Combinatorial models for the connected covering spaces have also been constructed, see for example [7, 19].

The crux of this paper is the generalization of Salvetti's result. We prove that the incidence relations in the face poset of a submanifold arrangement determine a cell complex that has the homotopy type of the tangent bundle complement (Theorem 3.7).

The paper is organized as follows. We state necessary background material in Section 1. In Section 2 we define *the arrangement of submanifolds* and present some examples. We then introduce the tangent bundle complement in Section 3 and prove that it has the homotopy type of a finite dimensional simplicial complex. We show that this simplicial complex is determined by the combinatorics of the incidence relations induced by submanifold intersections. In Section 4.1 we construct a regular CW complex which also has the homotopy type of this complement. This particular cell structure helps us better understand the relationships between the combinatorics and the topology in the context of submanifold arrangements. In Section 4.2 we show that this CW complex has a special combinatorial structure similar to that of zonotopes. We use the theory of *metrical-hemisphere complexes* (first introduced in [22]) to explain this combinatorial structure. Finally in Section 5 we study some homotopy theoretic aspects of the tangent bundle complement. First, we describe how the relations in the fundamental group of the complement depend on the face poset of the arrangement and also comment on the higher homotopy groups of the complement. We end with a description and characterization of the connected covering spaces in Section 5.2.

This particular generalization of hyperplane arrangements and the complexified complement is due to Tduesz Januskeweisz and Richard Scott. A video recording of Richard Scott's talk at M.S.R.I. is available at [23].

Acknowledgements. This paper is a part of the author's doctoral thesis [9]. The author would like to thank his supervisor Graham Denham. This work would not have been possible without the discussions with Tduesz Januskeweisz. Sincere thanks to Mario Salvetti for being the external examiner of the thesis and also for the warm hospitality in Pisa. Finally, the author would like to acknowledge the support of the Mathematics department at Northeastern University.

1. Background

We start by briefly reviewing some basic facts about (complexified) hyperplane arrangements.

1.1. Hyperplane Arrangements.

Hyperplane arrangements arise naturally in geometric, algebraic and combinatorial instances. They occur in various settings such as finite dimensional projective or affine (vector) spaces defined over field of any characteristic. In this section we will formally define hyperplane arrangements and the combinatorial data associated with it in a setting that is most relevant to our work.

Definition 1.1. A real, central *arrangement of hyperplanes* is a collection $\mathcal{A} = \{H_1, \dots, H_k\}$ of finitely many codimension 1 subspaces (*hyperplanes*) in $\mathbb{R}^l$, $l \geq 1$.

If we allow $\mathcal{A}$ to contain affine hyperplanes (i.e., translates of codimension 1 subspaces) we call $\mathcal{A}$ an affine arrangement. However we will mostly consider central arrangements. Hence, an arrangement of hyperplanes will always mean central, unless otherwise stated.

An arrangement is *essential* if the intersection of all the hyperplanes is the origin. For an affine subspace X of $\mathbb{R}^l$, the *contraction* of X in $\mathcal{A}$ is given by the sub-arrangement $\mathcal{A}_X := \{H \in \mathcal{A} \mid X \subseteq H\}$. The hyperplanes of $\mathcal{A}$ induce a stratification (cellular decomposition) of $\mathbb{R}^l$, components of each stratum are called *faces* (or cells) of $\mathcal{A}$.

There are two posets associated with $\mathcal{A}$, namely, the intersection lattice and the face poset which contain important combinatorial information about the arrangement.

Definition 1.2. The *intersection lattice* $L(\mathcal{A})$ of $\mathcal{A}$ is defined as the set of all intersections of hyperplanes ordered by reverse inclusion. The join of two elements is defined as the intersection of corresponding subspaces and the meet is defined as the smallest subspace containing the union.

Note that for affine arrangements, set of all intersections only form a poset and not a lattice.

Definition 1.3. The *face poset* $\mathcal{F}(\mathcal{A})$ of $\mathcal{A}$ is the set of all faces ordered by inclusion: $F \leq G$ if and only if $F \subseteq \overline{G}$.

Codimension 0 faces are called *chambers*, the set of all chambers is denoted by $\mathcal{C}(\mathcal{A})$. A chamber is bounded if and only if it is a bounded subset of $\mathbb{R}^l$. Two chambers C and D are adjacent if and only if they have a common face.

By a complexified hyperplane arrangement we mean that an arrangement of hyperplanes in $\mathbb{C}^l$ such that the defining equation of each hyperplane is real. Hence to every hyperplane arrangement in $\mathbb{R}^l$ there corresponds an arrangement of hyperplanes in $\mathbb{C}^l$. An important topological space associated with a real hyperplane arrangement is:

Definition 1.4. Let $\mathcal{A}$ denote a hyperplane arrangement in $\mathbb{R}^l$. The *complexified complement* of such an arrangement is denoted by $M(\mathcal{A})$ and defined as

$$M(\mathcal{A}) := \mathbb{C}^l \setminus (\bigcup_{H \in \mathcal{A}} H_{\mathbb{C}})$$

where $H_{\mathbb{C}}$ is the hyperplane in $\mathbb{C}^l$ with the same defining equation as $H \in \mathcal{A}$.

Note that since $M(\mathcal{A})$ is of real codimension 2 in $\mathbb{C}^l$, it is connected. In fact it is an open submanifold of $\mathbb{C}^l$ with the homotopy type of a finite dimensional CW complex [17, Section 5.1].

1.2. **The Salvetti Complex.** There is a combinatorial description of the homotopy type of $M(\mathcal{A})$ introduced by Salvetti in [21]. By a combinatorial description we mean a construction, using the combinatorial data, of a regular CW-complex of dimension n which has the same homotopy type as that of $M(\mathcal{A})$. Note that this particular cell complex, which we denote by $Sal(\mathcal{A})$, is defined using the face poset and not the intersection lattice.

The k-cells of $Sal(\mathcal{A})$ are in one-to-one correspondence with the pairs $[F, C]$, where F is a codimension k face of the given arrangement and C is a chamber whose closure contains F. A cell labeled $[F_1, C_1]$ is contained in the boundary of another cell $[F_2, C_2]$ if and only if $F_1 \leq F_2$ in $\mathcal{F}(\mathcal{A})$ and C_1, C_2 are contained in the same chamber of (the arrangement) $\mathcal{A}_{F_1}$ (with the attaching maps being homeomorphisms). The seminal result of Salvetti is:

Theorem 1.5 (Salvetti [21]). *Let $\mathcal{A}$ be an arrangement of real hyperplanes and $M(\mathcal{A})$ be the complement of its complexification inside $\mathbb{C}^l$. Then there is an embedding of $Sal(\mathcal{A})$ into $M(\mathcal{A})$ moreover there is a natural map in the other direction which is a deformation retraction.*

As we are going to generalize this construction and extensively study its combinatorial and topological properties we skip a detailed discussion here. There is another proof of Salvetti's result by Paris in [19]. We would like to mention this result since we adopt a similar strategy to prove our main result (Theorem 3.7).

Theorem 1.6 (Paris [19]). *Let $\mathcal{A}$ be an arrangement of real hyperplanes and $M(\mathcal{A})$ be the complement of its complexification inside $\mathbb{C}^l$. Then for every pair in the set $\{(F, C) \in \mathcal{F}(\mathcal{A}) \times \mathcal{C}(\mathcal{A}) \mid F \leq C\}$ there corresponds an open set of $M(\mathcal{A})$ such that the union of all these sets is an open covering of $M(\mathcal{A})$. Moreover each of these open sets are contractible and so are their intersections and consequently (the geometric realization of) the nerve of this open covering has the homotopy type of $M(\mathcal{A})$.*

In general the above geometric realization is the barycentric subdivision of the CW complex $Sal(\mathcal{A})$. As a result, every d-chain in the nerve corresponds to a closed d-cell in $Sal(\mathcal{A})$, and vice versa.

1.3. **Topological Combinatorics.** The main theme underlying our work is to understand interactions between combinatorics and topology in a particular context. For the sake of completeness and also in order to fix notation we lay out some foundation in this section. We start by a quick review of order complexes, the Nerve lemma and regular cell complexes. We assume the reader's familiarity with abstract simplicial complexes and posets. The main reference is Kozlov's book [15].

At the end we state relevant results from homotopy theory applied to combinatorial settings. The approach we want to take here is more concrete; we would like to leave the complete generality of diagrams over arbitrary categories and focus on small, directed categories. The reference is the work of Welker, Ziegler and Živaljević on homotopy colimits of spaces [24].

A *standard n-simplex* is defined as the convex hull of standard unit basis in $\mathbb{R}^{n+1}$. The following construction (and its generalizations) will be used as a path to move from combinatorics to topology.

Definition 1.7. Given a finite abstract simplicial complex Δ, its *standard geometric realization* is the topological space obtained by taking the union of standard k-simplex in $\mathbb{R}^{|V|}$, for all simplices of dimension k. Any topological space that is homeomorphic to the standard realization of Δ is called the *geometric realization* of Δ, and is denoted by $|\Delta|$.

By convention a topological statement about an abstract simplicial complex is actually a statement about its geometric realization. For the sake of simplicity, in this paper, we will not differentiate between an abstract simplicial complex and its geometric realization. We will let the context decide. A simplicial map between two simplicial complexes induces a continuous map between the corresponding geometric realizations.

Definition 1.8. The *order complex* $\Delta(P)$ of a poset P is the abstract simplicial complex on vertex set P whose k-faces are the k-chains in P. Again, we will not differentiate between the order complex and its geometric realization.

A poset map between two posets induces a simplicial (or continuous) map between the corresponding order complexes. Given a simplicial complex Δ, the set of its faces ordered by inclusion forms a poset which is called as the *face poset*. The order complex of this face poset is called the *(first) barycentric subdivision* denoted by $sd(\Delta)$. The spaces $|\Delta|$ and $|sd(\Delta)|$ are homeomorphic. Thus from a topological viewpoint simplicial complexes and posets can be considered equivalent notions.

Next we consider the *Nerve Lemma*, which often helps simplify a given topological space for combinatorial applications. Recall that an *open covering* of a topological space is a collection of open subsets of the space such that their union is the space itself. The *nerve of an open covering* is a simplicial complex whose vertices correspond to the open sets and a set of $k+1$ vertices spans a k-simplex whenever the corresponding $k+1$ open sets have a nonempty intersection (we will refer to the geometric realization of a nerve as the nerve itself). In general the nerve need not reflect the topology of the ambient space, but the following result gives a useful condition when it does.

Theorem 1.9 (Theorem 15.21 [15])**.** *If $\mathcal{U}$ is a finite open cover of a topological space X such that every non empty intersection of open sets in $\mathcal{U}$ is contractible, then $X \simeq$ nerve $(\mathcal{U})$.*

We now mention some basic facts about the cell complexes that frequently occur in Topological Combinatorics. It is often the case that given some combinatorial data one would like to synthetically construct a topological space consistent with the data. The geometric simplicial complex is one such example. We consider a more general class of *regular cell complexes.* A subset e of a topological space X is called a *closed* (*open*) *k-cell* if it is homeomorphic to (the interior of) the standard k-ball in $\mathbb{R}^k$.

Definition 1.10. A *regular cell complex* $(X, \mathcal{C})$ is a pair consisting of a Hausdorff space X and a finite collection $\mathcal{C}$ of open cells in X such that

(1) $X = \bigcup_{e\in\mathcal{C}} e$,

(2) the boundary $\overline{e} \setminus e$ of each cell is a union of some members of $\mathcal{C}$.

Given a (regular) CW complex X the set consisting of its cells ordered by inclusion forms a poset. This poset is called as the face poset and it is denoted by $\mathcal{F}(X)$. We state a result concerning regular cell complexes which is important from the combinatorial view point.

Theorem 1.11. *Let $(X, \mathcal{C})$ be a regular cell complex and $\mathcal{F}(X)$ be its face poset. Then*

$$\Delta(\mathcal{F}(X)) \cong X.$$

Furthermore, this homeomorphism can be chosen so that it restricts to a homeomorphism between $\overline{e}$ and $\Delta(\mathcal{F}_{\leq e})$, for all $e \in \mathcal{C}$.

A *diagram of spaces* indexed over a poset is a covariant functor $\mathcal{D}\colon P \to \mathrm{Top}$ from a finite poset P to the category topological spaces and continuous maps (in this case $\mathcal{D}$ will be called a P-diagram). Note that a poset is a small category, with objects being elements of the poset and morphisms being defined from p to q if and only if $p \geq q$.

Definition 1.12. Consider a diagram $\mathcal{D}\colon P \to \mathrm{Top}$ over a poset P, then the homotopy colimit of $\mathcal{D}$ is defined as

$$\mathrm{hocolim}\mathcal{D} := \coprod_{x \in P} (\Delta(P_{\leq x}) \times \mathcal{D}(x))/\sim$$

where Δ denotes the order complex and $P_{\leq x} := \{y \in P | y \leq x\}$. The equivalence relation $\sim$ is defined by following identifications

$$\Delta(P_{\leq x}) \times \mathcal{D}(y) \hookrightarrow \Delta(P_{\leq y}) \times \mathcal{D}(y)$$

and

$$\Delta(P_{\leq x}) \times \mathcal{D}(y) \xrightarrow{id \times f_{yx}} \Delta(P_{\leq x}) \times \mathcal{D}(x)$$

where $y \geq x$ and $f_{yx}\colon \mathcal{D}(y) \to \mathcal{D}(x)$.

In its simplest form the next theorem provides sufficient conditions for a map between two posets to be a homotopy equivalence at the level of order complexes.

Theorem 1.13 (Quillen's theorem A [24])**.** *Let $f\colon P \to Q$ be a poset map such that $\Delta(f^{-1}(Q_{\geq q}))$ is contractible $\forall q$. If $\mathcal{D}$ is any Q-diagram and $f^*\mathcal{D}$ the corresponding (pull back) P-diagram then,*

$$\mathrm{hocolim}_P(f^*\mathcal{D}) \xrightarrow{\simeq} \mathrm{hocolim}_Q\mathcal{D}.$$

We now look at an explicit connection between the order complex and the homotopy colimit construction.

Definition 1.14. Let $\mathcal{P}$ be a poset. A *diagram of posets* is a diagram indexed over P that assigns a poset Q_p to each $p \in P$.

Given a diagram of posets $\mathcal{D}$ we construct a new poset, called as the poset limit of $\mathcal{D}$. It is denoted by $\mathrm{Plim}\mathcal{D}$ and defined as follows:

$$\mathrm{Plim}\mathcal{D} := \coprod_{p \in \mathcal{P}} \{p\} \times Q_p.$$

The order relations are:

$$(p, q) \geq (p', q') \Leftrightarrow \begin{cases} p \geq p' \text{ and} \\ f_{pp'}(q) \geq q'. \end{cases}$$

For a diagram of posets its homotopy colimit is indeed an order complex as the following lemma suggests.

Lemma 1.15 (Simplicial Model Lemma [24])**.** *Let $\mathcal{D}\colon P \to \mathrm{Top}$ be a diagram of posets. Then*

$$\mathrm{hocolim}\mathcal{D} \simeq \Delta(\mathrm{Plim}\mathcal{D}).$$

The above lemma is a special case of the Grothendieck construction, which is used to define the homotopy colimit of a diagram of categories.

2. Arrangements of Submanifolds

In this section we propose a generalization of hyperplane arrangements. In order to do this we isolate their important characteristics:

1. there are finitely many codimension 1 subspaces (with nonempty intersections) disconnecting the ambient space,
2. there is a stratification of the ambient space into contractible subsets,
3. (the geometric realization of) the face poset (of the above stratification) has the homotopy type of the ambient space.

Any reasonable generalization of hyperplane arrangements should posses these properties. Since smooth manifolds are locally Euclidean they are obvious candidates for the ambient space. In this setting we can study arrangements of codimension 1 submanifolds that satisfy certain nice conditions, for example, locally, we would like our submanifolds to behave like hyperplanes.

We recall some relevant results from topological embeddings (the main reference is [20]). An l-dimensional (topological) manifold is a separable metric space in which each point has a *coordinate neighborhood* homeomorphic to $\mathbb{R}^l$. An *embedding* of a topological space X into a topological space Y is a homeomorphism $f\colon X \to Y$ of X onto a subset of Y. Two such embeddings f, g are said to be *equivalent* if there is a self homeomorphism h of Y such that $hf = g$. Our focus is on the codimension 1 smooth submanifolds that are *embedded as closed subsets*. These type of submanifolds behave very much like hyperplanes.

Lemma 2.1. *If X is a connected l-manifold and N is a connected $(l-1)$-manifold embedded in X as a closed subset, then $X \setminus N$ has either 1 or 2 components. If in addition $H_1(X, \mathbb{Z}_2) = 0$ then $X \setminus N$ has exactly two components (i.e. N separates X).*

Proof. The lemma follows from the following exact sequence of pairs in mod 2 homology:

$$H_1(X, \mathbb{Z}_2) \to H_1(X, X \setminus N, \mathbb{Z}_2) \to \tilde{H}_0(X \setminus N, \mathbb{Z}_2) \to \tilde{H}_0(X, \mathbb{Z}_2) = 0. \qquad \square$$

Lemma 2.2. *Let X be a connected l-manifold and let N be a connected, $(l-1)$-submanifold embedded in X as a closed subset. Then N separates X if and only if the inclusion induced homomorphism $H_c^{n-1}(X, \mathbb{Z}_2) \to H_c^{n-1}(N, \mathbb{Z}_2)$ (on the cohomology with compact supports) is trivial.*

Proof. The proof is a simple diagram chase:

$$\begin{tikzcd}
H_c^{n-1}(X, \mathbb{Z}_2) \arrow[r] \arrow[d, "\cong"'] & H_c^{n-1}(N, \mathbb{Z}_2) \cong \mathbb{Z}_2 \arrow[d, "\cong"] & & \\
H_1(X, \mathbb{Z}_2) \arrow[r] & H_1(X, X \setminus N, \mathbb{Z}_2) \arrow[r] & \tilde{H}_0(X \setminus N, \mathbb{Z}_2) \arrow[r] & 0.
\end{tikzcd}$$

$\square$

However in general it is not true that all codimension 1 submanifolds separate the ambient manifold there are weaker conditions.

Definition 2.3. A connected codimension 1 submanifold N in X is said to be *two sided* if N has a neighborhood U_N such that $U_N \setminus N$ has two connected components; otherwise N is said to be *one sided*. In general a disconnected codimension 1 submanifold is two sided if each of its connected component is two sided.

Note that being two sided is in some sense a local condition. For example, a point in S^1 does not separate S^1, however, it is two sided. The following corollary follows from the definition.

Corollary 2.4. *If X is a l-manifold and N is a $(l-1)$ manifold embedded in X as a closed subset, where $H_1(N, \mathbb{Z}_2) \cong 0$ then N is two sided.*

According to the above result $\mathbb{R}P^2 \hookrightarrow \mathbb{R}P^3$ is clearly not two sided. On the other hand there is a still weaker notion of separation which is always true.

Corollary 2.5. *Every codimension 1 submanifold N is locally two sided in X; that is, each $x \in N$ has arbitrarily small connected neighborhoods U_x such that $U_x \setminus (U_x \cap N)$ has two components.*

An n-manifold N contained in the interior of an l-manifold X is *locally flat* at $x \in N^\circ$, if there exists a neighborhood U_x of x in X such that $(U_x, U_x \cap N) \cong (\mathbb{R}^l, \mathbb{R}^n)$. An embedding $f\colon N \to X$ such that $f(N^\circ) \subset X^\circ$ is said to be *locally flat at a point* $x \in N$ if $f(N)$ is locally flat at $f(x)$. Embeddings and submanifolds are *locally flat* if they are locally flat at every point.

It is necessary to consider the locally flat class of submanifolds otherwise one could run into pathological situations. For example, the Alexander horned sphere, it is an (non flat) embedding of S^2 inside S^3 such that the connected components of its complement are not even simply connected (see [20, Page 65]). Finally we add a technical condition so that these locally flat submanifolds intersect like hyperplanes.

Definition 2.6. Let X be a manifold of dimension l and let $\{N_1, \dots, N_k\}, (k \geq 2)$ be codimension 1, locally flat submanifolds of X. We say that these submanifolds *intersect like hyperplanes* if and only if for every nonempty $Y \subseteq X$ which can be written as $Y = \cap_{i=1}^{j} N_i, (j \leq l)$ and for every $x \in Y$ there exists an open neighborhood V_x of x and a coordinate chart $\phi_x \colon V_x \to \mathbb{R}^l$ such that for each N_i containing x, the image $\phi_x(N_i \cap V_x)$ is a hyperplane passing through the origin.

See [9, Chapter 6] for an example of locally flat embeddings of planes in $\mathbb{R}^3$ that do not intersect like planes. The desired generalization of hyperplane arrangements is the following:

Definition 2.7. Let X be a connected, smooth, real manifold of dimension l. An **arrangement of submanifolds** is a finite collection $\mathcal{A} = \{N_1, \dots, N_k\}$ of codimension 1 locally flat smooth submanifolds in X such that

1. The N_i's intersect like hyperplanes.
2. $X \setminus N_i$ has exactly two connected components for every i.
3. The intersections of N_i's define a regular CW decomposition of X.

The submanifolds in an arrangement are allowed to have more than one connected component.

Before we convince the reader that this is the right setting, let us first look at how we can associate combinatorial data to such an arrangement and a few examples. Just like in the case of hyperplane arrangements, the intersection sets determined by these submanifolds have a combinatorial structure.

Definition 2.8. The *intersection poset* denoted by $L(\mathcal{A})$ is the set of connected components of all possible intersections of N_i's ordered by reverse inclusion, by convention $X \in L(\mathcal{A})$ as the smallest element. The rank of each element in $L(\mathcal{A})$ is defined to be the codimension of the corresponding intersection.

Definition 2.9. The intersections of these N_i's in $\mathcal{A}$ define a stratification of X. The connected components in each stratum are called *faces*. The set of all the faces, denoted by $\mathcal{F}(\mathcal{A})$, is the *face poset* with the ordering $F \leq G \Leftrightarrow F \subseteq \overline{G}$. The rank of each face is its dimension. Top dimensional faces are called chambers and the set of all chambers is denoted by $\mathcal{C}(\mathcal{A})$.

If there is no confusion about the arrangement, we will omit $\mathcal{A}$ and denote the two posets by $\mathcal{F}$ and L. The geometric realization of the face poset of an arrangement has the homotopy type of the ambient manifold (follows from the definition and Theorem 1.11).

The obvious examples of these submanifold arrangements are the hyperplane arrangements. Let us look at different types of arrangements.

Example 2.10. Let X be the circle S^1, a smooth one dimensional manifold, the codimension 1 submanifolds are points in S^1. Consider the arrangement $\mathcal{A} = \{p, q\}$ of a 0-sphere. Figure 1 shows this arrangement and the Hasse diagrams of the face poset and the intersection poset.



Figure 1. Arrangement of 2 points in a circle.

Example 2.11. As a 2-dimensional example consider an arrangement of 2 great circles N_1, N_2 in S^2. Figure 2 shows this arrangement and the related posets. The face poset has two 0-cells, four 1-cells and four 2-cells. Also note that the order complex of the face poset has the homotopy type of S^2.

Recently there has been an increase in interest to study arrangements in a torus (see for example [16]). We refer the reader to [9, Chapter 4] for an explicit treatment of toric arrangements, arrangements of spheres and of projective spaces. A few notations before we show that arrangements of submanifolds are locally arrangements of hyperplanes.



FIGURE 2. Arrangement of 2 circles in a sphere.

Definition 2.12. Let $\mathcal{A}$ be an arrangement of submanifolds, let $\mathcal{A}_{cc}$ denote the set of all connected components of members of $\mathcal{A}$. For a point $x \in X$ define the *local arrangement* at x as follows:

$$\mathcal{A}_x := \{N \in \mathcal{A}_{cc} \mid x \in N\}$$

Similarly for a face F local arrangement at F is:

$$\mathcal{A}_F := \{N \in \mathcal{A}_{cc} \mid F \subset N\}$$

The *restriction* of a local arrangement to an open set $V \subseteq X$ is:

$$\mathcal{A}_F|_V := \{N \cap V | N \in \mathcal{A}_F\}$$

Lemma 2.13. *Let F be a face of an arrangement $\mathcal{A}$ (that is $F \in \mathcal{F}(\mathcal{A})$). Then there exists an open set $V_F \subsetneq X$ containing F and a map $\phi\colon V_F \to \mathbb{R}^l$ such that*

1. *$V_F \cap N = \emptyset$ for every $N \notin \mathcal{A}_F$.*
2. *ϕ is a homeomorphism.*
3. *ϕ maps $\mathcal{A}_F|_{V_F}$ to a central arrangement of hyperplanes in $\mathbb{R}^l$.*

Proof. Each submanifold $N \in \mathcal{A}_{cc}$ has an open neighborhood in X which is homeomorphic to $N \times (-1, 1)$ (this follows from Brown's bicollared theorem [20, Theorem 1.7.5]). We call such an open neighborhood the bicollar of the submanifold. A face F is homeomorphic to an open cell of some dimension and is an intersection of finitely many codimension 1 submanifolds. Take the open set V_F to be the intersection of all the bicollars containing F. Moreover this V_F can be chosen such that it intersects with only those faces which contain $\overline{F}$.

Since X is a smooth manifold, there exists a homeomorphism $\phi\colon V_F \to \mathbb{R}^l$, say a coordinate chart. If necessary ϕ can be composed with a homeomorphism of $\mathbb{R}^l$ so that for every N containing F, the set $N \cap V_F$ is mapped to a codimension 1 subspace of $\mathbb{R}^l$. As ϕ is a homeomorphism it preserves the incidence relations between the faces, thus preserving the combinatorial structure of the local arrangement $\mathcal{A}_F|_{V_F}$. $\square$

Thus every point $x \in X$ has an open neighborhood V_x in X which is homeomorphic to a central arrangement of hyperplanes with x at its origin. The hyperplanes in that arrangement correspond to $N \cap V_x$ for every $N \in \mathcal{A}_x$.

3. The Tangent Bundle Complement

The aim of this section is to associate a connected topological space to submanifold arrangements that generalizes the construction of complexified hyperplane complement. If one were to forget the complex structure on $\mathbb{C}^l$ then, topologically, it is the tangent bundle of $\mathbb{R}^l$. Same is true for a hyperplane H and its complexification $H_{\mathbb{C}}$. Hence the complexified complement of a hyperplane arrangement can also be considered as a complement inside the tangent bundle. We use this topological viewpoint to define a generalization of the complexified complement for submanifold arrangements.

Definition 3.1. Let X be a l-dimensional manifold and $\mathcal{A} = \{N_1, \dots, N_k\}$ be an arrangement of submanifolds. Let TX denote the tangent bundle of X and let $T\mathcal{A} := \bigcup_{i=1}^{k} TN_i$. The **tangent bundle complement** of the arrangement $\mathcal{A}$ is defined as

$$M(\mathcal{A}) := TX \setminus T\mathcal{A}.$$

It is connected as it is of codimension 2 in TX.

For a point $x \in X$ and for all $N \in \mathcal{A}_x$, let $T_x(\mathcal{A}) = \bigcup T_x(N)$. Define the *tangent space complement* at $x \in X$ as

$$M(\mathcal{A}_x) := T_x(X) \setminus T_x(\mathcal{A}).$$

Then $M(\mathcal{A})$ can be rewritten as follows

$$M(\mathcal{A}) = \{(x, v) \mid x \in X, v \in M(\mathcal{A}_x)\}.$$

We start the study of $M(\mathcal{A})$ by first understanding the tangent space complement $M(\mathcal{A}_x)$.

Lemma 3.2. *Let $F \in \mathcal{F}(\mathcal{A})$ and let $\phi\colon V_F \to \mathbb{R}^l$ be a coordinate chart for an open neighborhood V_F of F. Then for every $x \in V_F$,*

$$M(\mathcal{A}_x) \cong \phi(V_F \setminus (\mathcal{A}_x|_{V_F})).$$

Proof. Observe that the linear map $(d\phi)_x\colon T_x(X) \to \phi(V_F)$ ($\cong \mathbb{R}^l$) is an isomorphism and for the same reasons $T_x(N) \cong \phi(N \cap V_F)$ ($\cong \mathbb{R}^{l-1}$) for every $N \in \mathcal{A}_x$. The map $\Phi\colon M(\mathcal{A}_x) \to \phi(V_F \setminus (\mathcal{A}_x|_{V_F}))$ defined by sending $(x, v) \mapsto (d\phi)_x((x, v))$ is a homeomorphism independent of the choice of coordinates. $\square$

Remark 3.3. Since $\phi(\mathcal{A}_x|_{V_F})$ is an arrangement of hyperplanes in $\phi(V_F)$ (Lemma 2.13), $T_x(\mathcal{A})$ is an arrangement of hyperplanes in $T_x(X)$. Moreover, the two arrangements have isomorphic face posets as well as intersection lattices.

As a consequence of the above lemma the connected components of the tangent space complement $M(\mathcal{A}_x)$ can be indexed by the chambers of $\mathcal{A}_x|_{V_F}$. Let F be a face and C be a chamber of $\mathcal{A}$ such that $F \subseteq \overline{C}$, for every $x \in F$ define

$$C_F(x) := \{(x, v) \in M(\mathcal{A}) \mid \Phi((x, v)) \in \phi(C \cap V_F)\}. \tag{3.1}$$

Figure 3 is a 2-dimensional example that illustrates Lemma 3.2, showing the two homeomorphisms and a component $C_F(x)$ of the tangent space complement.



FIGURE 3. The tangent space complement

Before stating the main theorem of this section let us mention one more definition. A submanifold N of $\mathcal{A}$ *separates* two chambers C and D if and only if they are contained in the distinct connected components of $X \setminus N$. For two chambers C, D the set of all the submanifolds that separate these two chambers is denoted by $R(C, D)$. The following lemma is now evident.

Lemma 3.4. *Let X be a l-manifold and $\mathcal{A}$ be an arrangement of submanifolds, let C_1, C_2, C_3 be three chambers of this arrangement. Then,*

$$R(C_1, C_3) = [R(C_1, C_2) \setminus R(C_2, C_3)] \bigcup [R(C_2, C_3) \setminus R(C_2, C_1)].$$

The *distance between two chambers* is defined as the cardinality of $R(C, D)$ and denoted by $d(C, D)$. Given a face F and a chamber C of a submanifold arrangement $\mathcal{A}$, define the action of F on C as follows:

Definition 3.5. A face F acts on a chamber C to produce another chamber $F \circ C$ satisfying:

1. $F \subset \overline{F \circ C}$
2. $d(C, F \circ C) = \min \{d(C, C') \mid C' \in \mathcal{C}(\mathcal{A}), F \subset \overline{C'}\}$.

Lemma 3.6. *With the same notation as above, the chamber $F \circ C$ always exists and is unique.*

Proof. First, note that $F \circ C = F$ if and only if F itself is a chamber. Let $\text{codim} F \geq 1$ so, $F = N_1 \cap \cdots \cap N_r$. Let $\mathcal{A}' = \{N_1, \dots, N_r\}$, the collection $\mathcal{A}'$ need not be an arrangement of submanifolds. However, $\mathcal{A}'$ defines a stratification of the manifold and we will refer to the codimension 0 components of this stratification as chambers. There exists a unique chamber C' of $\mathcal{A}'$ which contains C. Then define $F \circ C$ to be the unique chamber of $\mathcal{A}$ that is contained in C' and whose closure contains F. □

If , $F' \geq F$ are two faces then $F' \circ (F \circ C) = F' \circ C$. Also, if $F \subseteq \overline{C}$ then $F \circ C = C$. Now we state the main theorem of this section.

Theorem 3.7. *Let $\mathcal{A}$ be an arrangement of submanifolds in an l-manifold X with $\mathcal{F}$ as its face poset and $M(\mathcal{A})$ as the associated tangent bundle complement. Then there exists a finite open cover of $M(\mathcal{A})$ such that each open set is indexed by a face F and a chamber C whose closure contains F. Moreover, each of these open sets is contractible and their intersections are contractible.*

Proof. For every face F fix an open set V_F that completely contains F and has non-empty intersections with only those faces whose closures contain F (this is possible because of Lemma 2.13). Observe that $\{V_F \mid F \in \mathcal{F}\}$ is an open cover of X. Let $h \colon V_F \times \mathbb{R}^l \to \pi^{-1}(V_F)$ denote the local trivialization, where $\pi \colon TX \to X$ is the projection map.

Choose C, a chamber whose closure contains F.

For every point $x \in V_F$ the map $h_x := h(x, -) \colon \mathbb{R}^l \to \pi^{-1}(x)$ is a linear isomorphism. Let y be an arbitrarily chosen point in F and $C_F(y)$ be the connected component of $M(\mathcal{A}_y)$ corresponding to $\phi(C \cap V_F)$. Denote the open subset $h_y^{-1}(C_F(y)) \subseteq \mathbb{R}^l$ by C_F and define

$$W(F, C) := h(V_F \times C_F) \subsetneq \pi^{-1}(V_F).$$

For the sake of notational simplicity we will identify $V_F \times C_F$ with its image. $W(F, C)$ is an open and contractible subset of $M(\mathcal{A})$. For a chamber C we have $W(C, C) = C \times \mathbb{R}^l$.

Now we show that these open sets cover $M(\mathcal{A})$. Let $(x, v) \in M(\mathcal{A})$ be an arbitrary point. Suppose that $x \in F$ for some face F. As $v \in M(\mathcal{A}_x)$, so $\phi^{-1}(\Phi(v)) \in C \cap V_F$, where C is some chamber whose closure contains F. Therefore $(x, v) \in W(F, C)$. The intersection of any two such open sets is given by

$$W(F, C) \cap W(F', C') = (V_F \cap V_{F'}) \times (C_F \cap C'_{F'})$$

which is clearly contractible. □

Example 3.8. Consider an arrangement $\mathcal{A} = \{0\}$ in $\mathbb{R}$, it divides the real line in two chambers $A = (-\infty, 0)$ and $B = (0, \infty)$. The tangent bundle complement of this arrangement is the punctured plane, since $\{0\}$ is the only point whose tangent space is disconnected. For $p = 0$ we take $V_p = (-1, 1)$ and for the chambers we take $V_A = A$ and $V_B = B$ as the coordinate neighborhoods. With this the open cover of $M(\mathcal{A})$ is given by setting:

1. $W(p, A) = \{(x, v) \mid x \in V_p, v < 0\}$,
2. $W(p, B) = \{(x, v) \mid x \in V_p, v > 0\}$,
3. $W(A, A) = \{(x, v) \mid x \in A, v \in \mathbb{R}\}$,
4. $W(B, B) = \{(x, v) \mid x \in B, v \in \mathbb{R}\}$.

Figure 4 shows the open cover defined above of the tangent bundle complement.



FIGURE 4. Example of an open cover of the tangent bundle complement

The open covering $\{W(F,C) \mid F \leq C\}$ of $M(\mathcal{A})$ constructed in Theorem 3.7 satisfies the hypothesis of the Nerve Lemma. Hence the nerve of this open cover has the homotopy type of $M(\mathcal{A})$. In order to identify the simplices of this nerve we need to understand how these open sets intersect.

Lemma 3.9. *$W(F,C) \cap W(F',C') \neq \emptyset$ if and only if $F \leq F'$ and $F' \circ C = C'$.*

Proof. By construction of these open sets we have,

$$W(F,C) \cap W(F',C') = (V_F \cap V_{F'}) \times (C_F \cap C'_{F'})$$

Clearly $V_F \cap V_{F'} \neq \emptyset$ if and only if $F \leq F'$. We also need the other intersection to be nonempty,

$$\begin{aligned} C_F \cap C'_{F'} \neq \emptyset \quad &\Longleftrightarrow \quad (C \cap V_F) \cap (C' \cap V_{F'}) \neq \emptyset \ \text{ due to (3.1)} \\ &\Longleftrightarrow \quad (V_F \cap V_{F'}) \cap (C \cap C') \neq \emptyset \\ &\Longleftrightarrow \quad C \cap C' \neq \emptyset \\ &\Longleftrightarrow \quad \text{either } C' = F' \circ C \text{ (or } C' = C). \quad \square \end{aligned}$$

Let $\mathcal{S}$ denote an abstract set in bijection with the open sets in the above open cover. We denote by (F,C) the element of $\mathcal{S}$ corresponding to $W(F,C)$.

Lemma 3.10. *The relation $(F_2, C_2) \leq_s (F_1, C_1)$ if and only if $F_1 \leq F_2$ and $F_2 \circ C_1 = C_2$. defines a partial order on $\mathcal{S}$.*

Proof. The arguments are similar to the proof of [19, Lemma 3.1]. It is obvious that the relation is reflexive and symmetric, let us check the transitivity. Pick 3 elements such that $(F_3, C_3) \leq_s (F_2, C_2)$ and $(F_2, C_2) \leq_s (F_1, C_1)$.

The first inequality implies that $F_2 \leq F_3$ and $F_3 \circ C_2 = C_3$. Similarly from the second inequality we have, $F_1 \leq F_2$ and $F_2 \circ C_1 = C_2$. Since $\mathcal{F}$ is a poset, $F_3 \leq F_1$ and $C_3 = F_3 \circ (F_2 \circ C_1) = F_3 \circ C_1$ which concludes the transitivity. □

With this partial order we can now characterize the chains in the nerve.

Lemma 3.11. *Let $\mathcal{A}$ be an arrangement of submanifolds, then for a chain in $\mathcal{S}$ there corresponds a chamber and a chain in $\mathcal{F}(\mathcal{A})$.*

Proof. Let $F_0 \leq \cdots \leq F_k$ be a chain in $\mathcal{F}$ and let C be a chamber. Then the Lemma 3.10 implies that $(F_k, F_k \circ C) \leq_s \cdots \leq_s (F_0, F_0 \circ C)$ is a chain in $\mathcal{S}$. Moreover, using the same lemma it can be shown that every chain in $\mathcal{S}$ is of this form. □

Now we have proved that $\mathcal{S}$ is a partially ordered set and it follows that the geometric realization of $(\mathcal{S}, \leq_s)$ and the nerve described by the open sets $W(F, C)$ are homeomorphic. A k-simplex in this nerve is a k-chain in $(\mathcal{S}, \leq_s)$. Let $F_0 \leq \cdots \leq F_k$ be a chain in $\mathcal{F}(\mathcal{A})$ and let C be a chamber (such that $F_k \leq C$) then both of them determine a simplex in the nerve given by

$$(F_k, C) \leq_s \cdots \leq_s (F_0, C)$$

In fact every simplex of the nerve is of this form. We summarize all this discussion in the following definition.

Definition 3.12 (**The Salvetti Complex**)**.** Let X be a smooth l-manifold and $\mathcal{A}$ be an arrangement of codimension 1 submanifolds. Define the **Salvetti poset** as

$$\mathcal{S} = \{(F, C) \in \mathcal{F}(\mathcal{A}) \times \mathcal{C}(\mathcal{A}) \mid F \leq C\}$$

and the partial order as

$$(F, C) \leq_s (F', C') \text{ if and only if } F \leq F' \text{ and } F' \circ C = C'$$

The **Salvetti complex** $Sal(\mathcal{A})$ is defined as the geometric realization of $(\mathcal{S}, \leq_s)$ and it has the homotopy type of the tangent bundle complement $M(\mathcal{A})$.

Example 3.13. Let $\mathcal{A}$ be arrangement of 2 points in a circle. The complement $M(\mathcal{A})$ in this case is $TS^1 \setminus \{p, q\}$ (tangent bundle of a point is the point itself). As $TS^1 \cong S^1 \times \mathbb{R}$, the space $M(\mathcal{A})$ is an infinite cylinder with 2 punctures. Hence $M(\mathcal{A}) \simeq S^1 \vee S^1 \vee S^1$. Figure 5 shows the *Salvetti complex* for this arrangement.

The following theorem states the connection between the manifold X and the associated tangent bundle complement.

Theorem 3.14. *Let X be a smooth manifold and $\mathcal{A}$ be an arrangement of submanifolds, let $M(\mathcal{A})$ denote the associated tangent bundle complement. Then the manifold X is a retract of $M(\mathcal{A})$.*

Proof. Let $r\colon \mathcal{S} \to \mathcal{F}(\mathcal{A})$ be the map defined by $(F, C) \mapsto F$. This map is order reversing and induces a simplicial map on the geometric realizations which we will again denote by r. For a chamber C define a new map $\iota_C\colon \mathcal{F}(\mathcal{A}) \hookrightarrow \mathcal{S}$ by sending F to $(F, F \circ C)$. This map is injective and order reversing, consequently it induces a continuous injective map on the geometric realizations.



Figure 5. Salvetti complex

Let $X(C) \subset Sal(\mathcal{A})$ denote the image of ι_C, then $X(C) \cong X$ and in fact

$$Sal(\mathcal{A}) = \bigcup_{C \in \mathcal{C}(\mathcal{A})} X(C).$$

The composition $r \circ \iota_C$ is the identity on $|\mathcal{F}(\mathcal{A})|$ which establishes the claim. □

Intuitively speaking, a submanifold arrangement is made by gluing central hyperplane arrangements in a compatible way. It is natural to ask whether the associated tangent bundle complement is also made up of gluing spaces. We show that the complement $M(\mathcal{A})$ is a homotopy colimit of a diagram of spaces over the face poset. We make use of the fact that there is a combinatorial description of the complement. For the local arrangement $\mathcal{A}_F|V_F$ let $\mathcal{S}(\mathcal{A}_F)$ denote the associated Salvetti poset and $Sal(\mathcal{A}_F)$ denote its geometric realization.

Proposition 3.15. *Let X be a l-manifold and let $\mathcal{A}$ be an arrangement of codimension* 1 *submanifolds with $\mathcal{F}(\mathcal{A})$ as its face poset. If $F' \geq F$ are two faces then there is an inclusion $Sal(\mathcal{A}_{F'}) \hookrightarrow Sal(\mathcal{A}_F)$ as simplicial complexes.*

Proof. It suffices to prove that $\mathcal{S}(\mathcal{A}_{F'})$ embeds into the Salvetti poset $\mathcal{S}(\mathcal{A}_F)$. This follows if the face poset of $\mathcal{A}_{F'}|_{V_{F'}}$ includes into the face poset of $\mathcal{A}_F|_{V_F}$. But this is clear since $\mathcal{F}(\mathcal{A})_{\geq F'} \hookrightarrow \mathcal{F}(\mathcal{A})_{\geq F}$. □

Theorem 3.16. *Consider the diagram of spaces $\mathcal{D}\colon \mathcal{F}(\mathcal{A}) \to \mathrm{Top}$, given by setting $\mathcal{D}(F) = Sal(\mathcal{A}_F)$. The morphisms $\mathcal{D}(F' \to F)$ are the inclusions $Sal(\mathcal{A}_{F'}) \hookrightarrow Sal(\mathcal{A}_F)$. Then* $\mathrm{hocolim}\mathcal{D} \simeq Sal(\mathcal{A})$.

Proof. Let $\mathcal{E}$ be a diagram of posets indexed over $\mathcal{F}(\mathcal{A})$ such that it assigns $\mathcal{S}(\mathcal{A}_F)$ to every F. It follows from the simplicial model lemma (Lemma 1.15) that $\mathrm{hocolim}\mathcal{D} \simeq \Delta(\mathrm{Plim}\mathcal{E})$. Hence we need to show that the poset map $\phi\colon \mathrm{Plim}\mathcal{E} \to \mathcal{S}$ defined by $(F, (F', C)) \mapsto (F', C)$ induces a homotopy equivalence $\Delta(\mathrm{Plim}\mathcal{E}) \simeq Sal(\mathcal{A})$, which follows from of Quillen's Theorem A (Theorem 1.13). □

4. COMBINATORICS OF THE CELL STRUCTURE

In this section we construct a CW complex that has the homotopy type of the tangent bundle complement. Inspired by the work of Deligne [6] on simplicial arrangements, Salvetti, in his seminal paper [21], proved that the face poset of a real hyperplane arrangement defines a CW complex embedded inside the complexified complement and that the complex is a strong deformation retract of the complement. We now show that the same construction generalizes.

4.1. **Salvetti CW-complex.** Let $\mathcal{A}$ be an arrangement of submanifolds in a l-manifold X and let $\mathcal{F}(\mathcal{A})$ denote the associated face poset. By $(X, \mathcal{F}(\mathcal{A}))$ we mean the regular cell structure of X induced by the arrangement. Recall that $\mathcal{F}^*(\mathcal{A})$ denotes the dual face poset, we denote by pair $(X, \mathcal{F}^*(\mathcal{A}))$ the *dual cell structure*. Every k-cell in $(X, \mathcal{F}(\mathcal{A}))$ corresponds to $(l-k)$-cell in $(X, \mathcal{F}^*(\mathcal{A}))$ for $0 \leq k \leq l$. The partial order on the cells of $\mathcal{F}^*(\mathcal{A})$ will be denoted by $\prec$.

For the sake of notational simplicity we will denote the dual cell complex by $X^*(\mathcal{A})$ (and by X^* if the context is clear). The symbols C, D will denote vertices of X^* and the symbol F^k will denote a k-cell dual to the codimension k-face F of $\mathcal{A}$. Note that a 0-cell C is a vertex of a k-cell F^k in X^* if and only if the closure $\overline{C}$ of the corresponding chamber contains the $(l-k)$-face F. The action of the faces on chambers that was introduced in Definition 3.5 is also valid for the dual cells. The symbol $F^k \circ C$ will denote the vertex of F^k which is dual to the unique chamber closest to C.

Given an arrangement in an l-manifold construct a regular CW complex $S(\mathcal{A})$ of dimension l as follows:
The 0-cells of $S(\mathcal{A})$ correspond to 0-cells of X^*, which we denote by the pairs $\langle C; C\rangle$.
For each 1-cell $F^1 \in X^*$ with vertices C_1, C_2, assign two homeomorphic copies of F^1 denoted by $\left\langle F^1; C_1\right\rangle, \left\langle F^1; C_2\right\rangle$. Attach these two 1-cells in $S(\mathcal{A})_0$ (the 0-skeleton) such that

$$\partial \left\langle F^1; C_i\right\rangle = \{\langle C_1; C_1\rangle, \langle C_2; C_2\rangle\}.$$

Orient the 1-cell $\left\langle F^1; C_i\right\rangle$ so that it begins at $\langle C_i; C_i\rangle$, to obtain an oriented 1-skeleton $S(\mathcal{A})_1$.

By induction assume that we have constructed the $(k-1)$-skeleton of $S(\mathcal{A})$, $1 \leq k-1 < l$. To each k-cell $F^k \in X^*$ and to each of its vertex C assign a k-cell $\left\langle F^k; C\right\rangle$ that is isomorphic to F^k. Let $\phi(F^k, C) \colon \left\langle F^k; C\right\rangle \to S(\mathcal{A})_{k-1}$ be the same characteristic map that identifies a $(k-1)$-cell $F^{k-1} \subset \partial F^k$ with the k-cell $\left\langle F^{k-1}; F^{k-1} \circ C\right\rangle \subset \partial \left\langle F^k; C\right\rangle$. Extend the map $\phi(F^k, C)$ to whole of $\left\langle F^k; C\right\rangle$ and use it as the attaching map, hence obtaining the k-skeleton. The boundary of every k-cell in given by

$$\partial \left\langle F^k; C\right\rangle = \bigcup_{F \prec F^k} \langle F; F \circ C\rangle . \tag{4.1}$$

Now we prove a theorem that justifies the construction of this cell complex.

Theorem 4.1. *The regular CW complex $S(\mathcal{A})$ constructed above has the homotopy type of the tangent bundle complement $M(\mathcal{A})$.*

Proof. First, note that the above construction implies that $S(\mathcal{A})$ is indeed a regular cell complex. Let $\mathcal{S}(\mathcal{A})$ denote the poset of cells of $S(\mathcal{A})$ ordered by inclusion. It is isomorphic to the Salvetti poset described in the Definition 3.12. Consequently the barycentric subdivision of $S(\mathcal{A})$ is isomorphic to the (simplicial) Salvetti complex $Sal(\mathcal{A})$, hence to every k-cell $\langle F; C\rangle$ in $S(\mathcal{A})$ there corresponds an ideal $\mathcal{S}(\mathcal{A})_{\leq_s [F,C]}$ (a triangulation of that k-cell). The claim now follows from the application of the Theorem 1.11. ☐

Remark 4.2. In order to simplify the notations and the arguments we will not distinguish between the simplicial and the CW versions of the Salvetti complex. We use the notation

$[F, C]$ to denote either a cell or its barycenter and hope that the context will make it clear. More importantly, henceforth we will assume that the tangent bundle complement is equipped with the above defined cell structure.

Remark 4.3. The original proof of Salvetti (that appeared in [21]) in the context of hyperplane arrangements is more direct. His idea is to first identify an open covering of the complexified complement (as we did in Theorem 3.7). Rather than considering the nerve of this open cover abstractly, he constructs a cell complex embedded inside the complexified complement. The final step is to show that there is a strong deformation retraction from the complement on to the cell complex. A similar approach could be used in case of a tangent bundle complement. However, a proof involving this idea is very technical since the construction of the nerve as an embedded simplicial complex could be fairly complicated. Also, such a proof might not shed any new light and this is the reason we have taken an approach to construct the regular CW complex abstractly and then show that its barycentric subdivision is isomorphic to the nerve of the open covering.

Example 4.4. As an example of this construction consider the arrangement of 2 points in a circle (Example 2.10). The Figure 6 below illustrates the dual cell structure induced by the arrangement and the associated Salvetti complex.



FIGURE 6. Arrangement in S^1 and the associated Salvetti complex

We now look at some obvious properties of the above defined CW structure and also infer some more information about the tangent bundle complement.

Theorem 4.5. *Let $\mathcal{A}$ be an arrangement of submanifolds in a l-manifold X and let $Sal(\mathcal{A})$ denote the associated cell complex. Then*

(1) *There is a natural cellular map $\psi\colon Sal(\mathcal{A}) \to X^*(\mathcal{A})$ given by $[F^k, C] \mapsto F^k$. The restriction of ψ to the 0-skeleton is a bijection and in general*
$$\psi^{-1}(F^k) = \{C \in \mathcal{C}(\mathcal{A}) | C \prec F^k\}$$

(2) *For every chamber C there is a cellular map $\iota_C\colon X^*(\mathcal{A}) \to Sal(\mathcal{A})$ taking F^k to $[F^k, F^k \circ C]$ which is an embedding of $X^*(\mathcal{A})$ into $Sal(\mathcal{A})$, and*
$$Sal(\mathcal{A}) = \bigcup_{C \in \mathcal{C}(\mathcal{A})} \iota_C(X^*).$$

(3) *The absolute value of the Euler characteristic of* $M(\mathcal{A})$ *is the number of bounded chambers.*

(4) *Let* $T\mathcal{A}$ *denote the union of the tangent bundles of the submanifolds in* $\mathcal{A}$ *then,*

$$\operatorname{rank} \tilde{H}^i(TX, T\mathcal{A}) = \begin{cases} |\chi(M(\mathcal{A}))| & \text{if } i = l \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Claims (1) and (2) follow from the construction and Theorem 3.14.

We prove (3) by explicitly counting cells in the Salvetti complex. The Euler characteristic of a CW complex K is equal to the alternating sum of number of cells of each dimension. Given a k-dimensional dual cell F^k there are as many as $|\{C \in \mathcal{C}(\mathcal{A}) | F \leq C\}|$ k-dimensional cells in $Sal(\mathcal{A})$. Hence for a vertex $[C, C] \in Sal(\mathcal{A})$ the number of k-dimensional cells that have this particular 0-cell as a vertex is equal to the number of k-faces of $\mathcal{F}^*(\mathcal{A})$ that contain C. The alternating sum of number of cells that contain a particular vertex C of $\mathcal{F}^*(\mathcal{A})$ is equal to $1 - \chi(Lk(C))$, where $Lk(C)$ is the link of C in $X^*(\mathcal{A})$. Applying this we get,

$$\chi(Sal(\mathcal{A})) = \sum_{C \in \mathcal{C}(\mathcal{A})} (1 - \chi(Lk(C)))$$

If a chamber is unbounded then $Lk(C) \simeq \mathbb{B}^l$ and on the other hand if it is bounded then $Lk(C) \simeq S^{l-1}$. Hence we have,

$$\begin{aligned} \chi(Sal(\mathcal{A})) &= \sum_{C \in \mathcal{C}(\mathcal{A})} (1 - \chi(Lk(C))) \\ &= \sum_{C \text{ unbounded}} (1 - 1) + \sum_{C \text{ bounded}} (1 - [1 + (-1)^{l-1}]) \\ &= (-1)^l \sum_{C \text{ bounded}} 1 \end{aligned}$$

Hence,

$$\chi(M(\mathcal{A})) = (-1)^l (\text{number of bounded chambers}).$$

Let $\bigcup \mathcal{A}$ denote the union of submanifolds in $\mathcal{A}$. Since $\mathcal{A}$ induces a regular cell decomposition it has the homotopy type of wedge of $(l-1)$-dimensional spheres. The number of spheres is equal to the number of bounded chambers. The claim (4) follows from the homeomorphism of pairs $(TX, T\mathcal{A}) \cong (X, \bigcup \mathcal{A})$. □

4.2. **Metrical-hemisphere complexes.** We now take a closer look at the cells of a Salvetti complex. Our aim is to understand how the combinatorial properties of the Salvetti complex associated to an arrangement of hyperplanes generalize in the context of submanifold arrangements. In particular, we show that even in the general setting the combinatorics of the cell structure is similar to that of a zonotope. Recall that a central arrangement of hyperplanes decomposes the ambient Euclidean space into open polyhedral cones. As a matter of fact every hyperplane arrangement is a normal fan of a very special polytope known as the zonotope. Zonotopes can be defined in various ways: for example, projections of cubes, Minkowski sums of line segments, dual (polar) of hyperplane arrangements etc. For more on the relationship between zonotopes and hyperplane arrangements see [25, Lecture 7].

Definition 4.6. A *zonotope* is a polytope all of whose faces are centrally symmetric (equivalently every 2-face is centrally symmetric). A *zonotopal cell* is a (closed) k-cell such that its face poset is isomorphic to the face poset of a k-zonotope for some k.

The face poset of a zonotope has some special combinatorial properties, the most important of which is the product structure. This product is basically the one on the face poset of a hyperplane arrangement or on the set of covectors of an oriented matroid. We show that the dual cells of a submanifold arrangement and the cells of its associated Salvetti complex enjoy similar combinatorial structure. In order to do this we do not use the language of zonotopes (since they are exclusive to Euclidean settings) but use the language of *metrical-hemisphere complexes.* These cell complexes possess all the essential combinatorial properties of a zonotope. The metrical-hemisphere complexes (MH-complexes for short) were first introduced in [22] where Salvetti generalized his construction and proved an analogue of Deligne's theorem for oriented matroids.

Let Q denote a connected, regular, CW complex (and $|Q|$ be the underlying space). The 1-skeleton of such a complex Q is a graph $G(Q)$ with no loops (abbreviated to G if the context is clear). The vertex set of this graph will be denoted by VG and the edge set by EG. An edge-path in $G(Q)$ is a sequence $\alpha = (l_1, \ldots, l_n)$ of edges that correspond to a connected path in $|Q|$. The inverse of a path is again a path $\alpha^{-1} = (l_n, \ldots, l_1)$. Two paths are composed by concatenation if ending vertex of one of the paths is the starting vertex of another. The distance $d(v, v')$ between two vertices will be the least of the lengths of paths joining v to v'. Given an i-cell $e^i \in Q$, $Q(e^i) := \{e^j \in Q : |e^j| \subset |e^i|\}$ and let $V(e^i) = VG \cap Q(e^i)$.

Definition 4.7. A regular CW complex Q is a QMH-complex (quasi-metrical-hemisphere complex) if and only if there exist two maps

$$\underline{\omega}, \overline{\omega} \colon VG \times Q \to VG$$

such that for all $v \in VG, e^i \in Q$ following properties are satisfied.

(1) $\underline{\omega}(v, e^i) \in V(e^i)$ and $d(v, \underline{\omega}(v, e^i)) = \text{minimum}\{d(v, u) | u \in V(e^i)\}$.
(2) $\overline{\omega}(v, e^i) \in V(e^i)$ and $d(v, \overline{\omega}(v, e^i)) = \text{maximum}\{d(v, u) | u \in V(e^i)\}$.
(3) $d(v, \overline{\omega}(v, e^i)) = d(v, u) + d(u, \overline{\omega}(v, e^i))$ for all $u \in V(e^i)$.

This definition imposes a strong restriction on the 1-skeletons of such complexes (see [22, Proposition 1]).

Lemma 4.8. *If Q is a QMH-complex then each circuit in G has an even number of edges.*

The next corollary follows from the above lemma and the definition of a zonotope.

Corollary 4.9. *Let Q be a closed k-cell which is a QMH-complex. Then Q is a zonotopal cell.*

For any $e^i \in Q$, indicate by $G(e^i) \subset G(Q)$ the subgraph corresponding to the 1-skeleton of e^i and by $d_{G(e^i)}$ the distance computed using $G(e^i)$.

Definition 4.10. A regular CW complex will be called a LMH-complex (local-metrical-hemisphere complex) if and only if each $Q(e^i)$ is a QMH-complex with respect to $d_{G(e^i)}$. Moreover, the following compatibility condition also holds: if $e^k \in Q(e^i) \cap Q(e^j), v \in V(e^i) \cap V(e^j)$ then

$$\underline{\omega}_{(e^j)}(v, e^k) = \underline{\omega}_{(e^i)}(v, e^k), \quad \overline{\omega}_{(e^j)}(v, e^k) = \overline{\omega}_{(e^i)}(v, e^k).$$

Here, $\underline{\omega}_{(e^j)}, \overline{\omega}_{(e^j)}$ are defined similar to $\underline{\omega}, \overline{\omega}$ but using $d_{G(e^j)}$.

Finally, Q will be called a **MH-complex** if Q is both a QMH-complex and a LMH-complex and for all $e^i \in Q, e^j \in Q(e^i), v \in V(e^i)$

$$\underline{\omega}(v, e^j) = \underline{\omega}_{(e^i)}(v, e^j), \quad \overline{\omega}(v, e^j) = \overline{\omega}_{(e^i)}(v, e^j).$$

Remark 4.11. Note that the 1-skeleton of a MH-complex has very special properties with respect to the distance. It is not enough to have a cell complex all of whose cells are zonotopal. Here are two examples that illustrate the special nature of MH-complexes.

Consider a 2-dimensional cell complex made up of two 1-cells and two 2-cells. The 1-cells are attached to an octagonal 2-cell. There is a one more trapezoidal 2-cell whose three 1-cells are attached to three 1-cells in the boundary of the octagonal cell as shown in Figure 7. The resulting complex is QMH but not LMH. Consider the 1-cell labeled by e in the figure. There are two vertices, namely v_1, v_2, in its boundary. Considering e as a member of the trapizoidal cell we see that the vertex v_1 is closest to the vertex v_4. On the other hand as a member of the octagonal 2-cell vertex v_2 is closest to v_4.



FIGURE 7. A QMH complex without LMH structure.

The next example shows a cell complex (Figure 8) obtained by removing the trapezoidal 2-cell from the first example. The resulting cell complex is both QMH and LMH but not a MH-complex. Consider the 1-cell labeled by e, there are two boundary vertices v_1, v_2. Considering e as a member of the octagonal cell the vertex v_2 is closest to v_3. But in the whole complex the boundary vertex of e closest to v_3 is v_1.



FIGURE 8. A QMH and LMH-complex which is not a MH-complex

The following lemma establishes the combinatorial connection between zonotopes and MH-complexes. It states that the distance between any two vertices is the same no matter how it is measured, either locally or globally (see [22, Proposition 5]).

Lemma 4.12. *Let Q be a MH-complex, $e^i \in Q, v, v' \in V(e^i)$. Then $d(v, v') = d_{G(e^i)}(v, v')$.*

Proof. Let $\alpha = (l_1, \dots, l_n)$ be a minimal path of $G(e^i)$ between v and v' (so $d_{G(e^i)}(v, v') = n$). Let v_{j-1}, v_j be the vertices of l_j ordered according to the orientation of α from v to v'. Since α is minimal in $G(e^i)$ and Q is a MH-complex we have,

$$\underline{\omega}_{(e^i)}(v, l_j) = v_{j-1} = \underline{\omega}(v, l_j)$$

Hence, $d_{G(e^i)}(v, v_j) = d_{G(e^i)}(v, v_{j-1}) + 1$ and $d(v, v_j) = d(v, v_{j-1}) + 1$ for $j = 1, \dots, n$. □

We now state the theorem that generalizes the relationship between hyperplane arrangements and zonotopes.

Theorem 4.13. *Let X be a smooth manifold of dimension l, $\mathcal{A}$ denote an arrangement of submanifolds. The (dual) cell complex $X^* = (X, \mathcal{F}^*)$ is a MH-complex.*

Proof. First, we need to define the two maps $\underline{\omega}, \overline{\omega}$ and then show that they are well defined. Let F^i be an i-cell and C be a vertex of X^* then,

$$\underline{\omega}(C, F^i) := F^i \circ C.$$

Using the same strategy as in the proof of Lemma 3.6 we can identify a unique chamber (of $\mathcal{A}$) whose closure contains (dual of) F^i and is farthest from C, denote this chamber by $F^i * C$ and

$$\overline{\omega}(C, F^i) := F^i * C.$$

This shows that the maps $\underline{\omega}, \overline{\omega}$ are well defined. Second, note that a path between two vertices C, C' has minimal length (among all paths from C to C') if and only if it crosses the faces that separate C from C' exactly once and does not cross any other face. The distance between any two vertices of X^* is thus

$$d(C, C') := |R(C, C')|.$$

Observe that if $F^i = N_1 \cap \cdots \cap N_r$ then $R(F^i \circ C, F^i * C) = \{N_1, \dots, N_r\}$ and if $D \in V(F^i)$ then

$$R(F^i \circ C, D) \bigcup R(F^i * C, D) = R(F^i \circ C, F^i * C) \text{ and } R(F^i * C, D) \cap R(F^i \circ C, D) = \emptyset$$

(by Lemma 3.4). Moreover,

$$R(C, D) = R(C, F^i \circ C) \bigcup R(F^i \circ C, D), \quad R(C, F^i \circ C) \cap R(F \circ C, D) = \emptyset$$

and

$$R(C, D) = R(C, F * C) \bigcup R(F * C, D), \quad R(C, F * C) \cap R(F * C, D) = \emptyset.$$

Using the last equality we see that X^* is a QMH-complex. The other compatibility conditions also follow easily. □

In a nutshell the above theorem justifies the following statement. MH-complexes are to submanifold arrangements as zonotopes are to hyperplane arrangements.

In case of hyperplane arrangements, each cell of the associated Salvetti complex is zonotopal (see [1, Proposition 5.7]). We generalize this claim and also show that in general the Salvetti complex is a MH-complex.

Theorem 4.14. *Let $\mathcal{A}$ be an arrangement of submanifolds in a manifold X and $Sal(\mathcal{A})$ denote the associated Salvetti complex. Then $Sal(\mathcal{A})$, with the CW structure described above, is a MH-complex.*

Proof. The 1-skeleton of $Sal(\mathcal{A})$ is obtained by 'doubling' the edges in the 1-skeleton of X^*. Hence the distance between any two vertices of $Sal(\mathcal{A})$ is same as the distance between the corresponding vertices of X^*. Also, by construction, there is a one-to-one correspondence between vertices of $[F^i, C]$ and the vertices of F^i for all $F^i \in X^*$. □

Remark 4.15. Note that in the construction of the Salvetti complex as mentioned before we used the $\underline{\omega}(F, C) = F \circ C$ map. It is possible to repeat the same construction using the $\overline{\omega}$ map. However these two complexes are isomorphic.

As a consequence of the Corollary 4.9, every 2-cell of the Salvetti complex (of an arrangement) is combinatorially equivalent to an oriented polygon with an even number of edges. This observation together with the condition (3) in Definition 4.7 (QMH-complex) proves the following.

Corollary 4.16. *In the Salvetti complex $Sal(\mathcal{A})$, the (oriented) 1-skeleton of a k-cell $[F^k, C]$ is composed of edge-paths going from $[C, C]$ to $[F^k * C, F^k * C]$. All these paths have same lengths and are directed away from $[C, C]$.*

In particular, the boundary of each 2-cell $[F^2, C]$ is composed of two equi-oriented edge-paths of the same length from $[C, C]$ to $[F^2 * C, F^2 * C]$. Moreover, the length of such a path is equal to the distance between its end vertices.

Proof. Assume that there is an edge $[F^1, C']$ contained in the boundary of $[F^k, C]$. Hence, $C' = F^1 \circ C$ and the result follows because of the condition (3) defining QMH-complex. □

We end this section with homotopy colimits. The next result follows from the observation that the barycentric subdivision of any regular cell complex is a poset limit (Definition 1.14).

Theorem 4.17. *Given an l-manifold X and a submanifold arrangement $\mathcal{A}$, let $\mathcal{F}^*$ denote the dual face poset. Define a diagram of spaces $\mathcal{D}\colon \mathcal{F}^* \to \mathrm{Top}$ with discrete spaces*

$$\mathcal{D}(F^i) := \{C \in X_0^* | C \prec F^i\}$$

and maps being the inclusions

$$\begin{array}{rcll} \mathcal{D}(F^i \to F^j)\colon \mathcal{D}(F^i) & \to & \mathcal{D}(F^j) & (i > j) \\ C & \mapsto & F^j \circ C & \end{array}$$

Then $\mathrm{hocolim}_{\mathcal{F}^*}\mathcal{D} \cong Sal(\mathcal{A})$.

Proof. Since all the space in the above diagram are in fact finite sets of points, we can consider these sets as posets with trivial order relation ($C \leq C' \Leftrightarrow C = C'$). In this situation we can apply the *simplicial model lemma* (Lemma 1.15). Hence the homotopy colimit of $\mathcal{D}$ has the homotopy type of $\Delta(\mathrm{Plim}\mathcal{D})$. The vertex set of $\Delta(\mathrm{Plim}\mathcal{D})$ is $\{(F, C) \in \mathcal{F}(\mathcal{A}) \times \mathcal{C}(\mathcal{A}) | F \subset \overline{C}\}$.

In order to determine higher dimensional simplices we look at the chains in $\mathrm{Plim}\mathcal{D}$. According to the Definition 1.14, the order relation on $\mathrm{Plim}\mathcal{D}$ is the following

$$(F, C) \leq (F', C') \Leftrightarrow F \prec F' \text{ in } \mathcal{F}^* \text{ and } F \circ C' = C.$$

Then it follows that the poset $\mathrm{Plim}\mathcal{D}$ is isomorphic to the Salvetti poset, which establishes the claim. □

5. HOMOTOPY THEORY OF THE TANGENT BUNDLE COMPLEMENT

In this section we focus on the homotopy theory of the tangent bundle complement. We start by defining the arrangement groupoid. This is a generalization of the groupoid *Gal* introduced by Deligne in [6] and also of the oriented systems introduced by Paris in [18]. We then look at the covering spaces of the Salvetti complex, our aim is to show that they are also determined combinatorially. We prove this by generalizing the theory of *Salvetti-type diagram models* introduced by Delucchi in [7] (building on the work of Paris [18]) to submanifold arrangements. The classification of the covering spaces is done at the end.

5.1. **The fundamental group(oid).** We start by extending the idea of an *arrangement graph*, associated with a hyperplane arrangement (see [6, 7, 18]).

Definition 5.1. Given an arrangement of submanifolds $\mathcal{A}$ in a manifold X, the arrangement graph, denoted by $G(\mathcal{A})$, is the oriented 1-skeleton of the associated Salvetti complex.

A *path* in the arrangement graph is a sequence of edges traversed not necessarily according to their orientations. The length of a path is equal to the number of edges traversed. An edge is a path of length 1 and vertex is path of length 0. Initial vertex of a path α is denoted by $s(\alpha)$ and the terminal vertex by $t(\alpha)$. Such directed paths can be represented by words whose letters correspond to edges. If α is an undirected path then $\alpha = (\epsilon_1 a_1, \dots, \epsilon_n a_n)$ where $\epsilon_i = \pm 1$ depending on whether a_i is traversed along or opposite to its orientation, the length of α is equal to 1 and $s(\epsilon_{i+1} a_{i+1}) = t(\epsilon_i a_i)$ for every i. If α and β are two paths such that $s(\beta) = t(\alpha)$ then their concatenation gives a new path denoted by $\alpha\beta$ (note that this is not a composition of functions). Inverse of a path $\alpha = (\epsilon_1 a_1, \dots, \epsilon_n a_n)$ is the path $\alpha^{-1} = (-\epsilon_n a_n, \dots, -\epsilon_1 a_1)$.

A *directed path* is a sequence of consecutive edges traversed according to their orientations. For example, $\alpha = (+a_1, \dots, +a_n)$ is a directed path of length n. A *minimal positive path* is a path having shortest length among all the positive paths that join its endpoints. Unless otherwise stated by a *path* we mean an *undirected path* and by a *positive path* we mean a directed path.

Recall that every 2-cell in the Salvetti complex is a polygon with an even number of (oriented) edges and that the boundary of such a 2-cell is composed of two minimal positive paths of same length (Corollary 4.16).

We define two types of '*moves*' on the paths in the arrangement graph:

1. If α is a path such that $\alpha = \alpha_1\gamma\alpha_2$ and if γ, γ' are two minimal positive paths joining the opposite vertices of a 2-cell, then substitute γ' for γ in α.
2. If α is a path such that $\alpha = \alpha_1\gamma\alpha_2$, where γ is a sub-path of a boundary path of a 2-cell of $Sal(\mathcal{A})$, substitute for γ the path γ' having same endpoints (i.e. $s(\gamma') = s(\gamma), t(\gamma') = t(\gamma)$) and containing all the remaining edges of that 2-cell.

Note that the above 'moves' define a homotopy between the directed paths.

Definition 5.2. Given a submanifold arrangement $\mathcal{A}$, the associated *category of positive paths*, (positive category for short) denoted by $\mathcal{G}^+(\mathcal{A})$, is the category of directed paths in the Salvetti complex.

Recall that (see, for example, [14, Chapter 3]) the objects of this category are the vertices of the arrangement graph. The morphisms are equivalence classes of positive paths. Two positive paths α and β from a vertex C to a vertex D are equivalent if and only if α is obtained from

β by a finite sequence of moves of type (1). It means that the two paths are connected by a sequence of substitutions of minimal positive paths. This equivalence relation will be called the *positive equivalence*, the equivalence class of a path α will be denoted by $[\alpha]_+$ and the relation denoted by $\overset{+}{\sim}$.

In case of hyperplane arrangements any two minimal positive paths with the same end points are homotopic in the associated Salvetti complex. Hence represent the same morphism in the positive category. However this need not be true for submanifold arrangements. For example, consider the Salvetti complex (Figure 6) associated to the arrangement of 2 points in S^1 (Example 2.10). The two edges labeled $[p, A]$ and $[q, A]$ are minimal positive paths with the same end points however they are not positive equivalent.

Lemma 5.3. *Let F be a face of a submanifold arrangement and C be a chamber such that $C \prec F$ in $\mathcal{F}^*$. Then any two minimal positive paths from C to $F * C$ are positive equivalent.*

Proof. The statement follows from the application of Corollary 4.16. Any two minimal positive paths from C to $F * C$ are in fact contained in the boundary of the cell $[F, C]$. □

Hence we can rephrase the positive equivalence as follows. Let α, α' be two positive paths from C to C'. Set $\alpha \overset{+}{\sim} \alpha'$ if and only if $\alpha = \alpha_1\beta\alpha_2$, $\alpha' = \alpha_1\beta'\alpha_2$ such that

(1) α_1 is a positive path from C to C'',
(2) β and β' are minimal positive paths from C'' to $F * C''$ (for some $C'' \prec C$),
(3) α_2 is a positive path from $F * C''$ to C'.

Definition 5.4. The *arrangement groupoid* of a submanifold arrangement is a category whose objects are the vertices of the arrangement graph. The morphisms in this category are equivalence classes of paths. Two paths are equivalent if and only if one is obtained from the other by a finite sequence of moves of type (2). The category will be denoted by $\mathcal{G}(\mathcal{A})$ and the morphisms will be denoted by $[\alpha]$, where α is a representative path.

The reader familier with homotopy theory will realize that the arrangement groupoid is the free groupoid on the Salvetti poset (consequently the fundamental groupoid of its nerve, the Salvetti complex). We state this as a result.

Lemma 5.5. *Let $\mathcal{A}$ be an arrangement of submanifolds in a manifold X. The arrangement groupoid $\mathcal{G}(\mathcal{A})$ is equivalent to the fundamental groupoid $\tilde{\pi}_1(Sal(\mathcal{A}))$ of the associated Salvetti complex.*

Before moving on we would like to explicitly state the relationship between the above defined categories.

Theorem 5.6. *For a submanifold arrangement $\mathcal{A}$ the category $\mathcal{G}(\mathcal{A})$ can be identified with the category of fractions of $\mathcal{G}^+(\mathcal{A})$ and the associated canonical functor $J\colon \mathcal{G}^+ \to \mathcal{G}$ is defined by sending $[\alpha]_+$ to $[\alpha]$.*

Proof. The proof follows from the observation that the moves of type (2) are composed of moves of type (1) and the definition of category of fractions (see [10, Chapter 1]). □

By studying the 2-skeleton of $Sal(\mathcal{A})$ we can deduce a presentation for the fundamental group.

Theorem 5.7. *Let X be a l-manifold and $\mathcal{A}$ be an arrangement of submanifolds with $Sal(\mathcal{A})$ being the associated Salvetti complex. If X is simply connected then two minimal positive paths in the* 1*-skeleton of $Sal(\mathcal{A})$ that have same initial as well as terminal vertex are homotopic.*

Proof. Let α, β be two positive paths such that $s(\alpha) = s(\beta) = [C, C]$, $t(\alpha) = t(\beta) = [D, D]$ and both of them have same lengths $(= d(C, D))$. Clearly, $\psi(\alpha), \psi(\beta) \subset X_1^*$ (recall that $\psi \colon Sal(\mathcal{A}) \to X^*$ was defined in Theorem 4.5) are minimal paths of length $d(C, D)$, so no two edges of these two paths are sent (by ψ) to the same edge of X^*. Without loss of generality, let $a = [F^1, C']$ be an edge of the path α, consider (a) as a sub-path of α. Let $s(a) = [C', C']$ then, $F^1 \circ C = C'$ since α is minimal. Repeating same argument for every edge of β we get that the paths α and β are contained in $\iota_C(X^*)$. Hence by Theorem 4.5 and the fact that $\pi_1(X^*) = \{1\}$, it follows that α and β are homotopic. ☐

An immediate consequence of this theorem is the following:

Corollary 5.8. *If X is a l-manifold, $\mathcal{A}$ is an arrangement of submanifolds and $M(\mathcal{A})$ is the tangent bundle complement then,*

$$\pi_1(M(\mathcal{A})) \cong \pi_1(G(\mathcal{A}))/\{\alpha\beta^{-1} \mid \alpha, \beta \ \text{ are minimal positive paths with same end points}\}.$$

Proof. A loop in the arrangement graph $G(\mathcal{A})$ is contractible if and only if it is the boundary of a 2-cell (of $Sal(\mathcal{A})$). According to Corollary 4.16 the boundary of every 2-cell is composed of two minimal positive paths with same ends. ☐

In particular Theorem 5.7 applies to hyperplane arrangements and arrangements in a sphere (dimension ≥ 2). Minimal positive paths with same end points are homotopic in the Salvetti complex. Hence these paths represent the same morphism in the corresponding arrangement groupoid. More is true for (central) hyperplane arrangements; minimal positive paths with same end points represent same morphism in the positive category (see [6, 1.12] for the original proof). This means that the restriction of functor J to the class of minimal positive paths is faithful (this holds true even for arrangements of pseudospheres [22, Theorem 20] and pseudohyperplanes [9, Chapter 5]).

Lemma 5.9. *Let $\mathcal{A}$ be an arrangement of submanifolds in a l-manifold X. If $\pi_n(X) \neq 0$ then $\pi_n(M(\mathcal{A})) \neq 0$.*

Proof. Follows from Theorem 3.14. ☐

The only hope to find $K(\pi, 1)$ complements is to consider arrangements in $K(\pi, 1)$ manifolds.

5.2. **Salvetti-type diagram models.**

The coveing spaces of a topological space can be classified by their fundamental groupoids. According to [2, Chapter 10], the connected covers of the fundamental groupoid correspond to fundamental groupoids of the corresponding covering spaces (see also [14, Chapter 13]). The fundamental group of the Salvetti complex is isomorphic to the object group $\pi(\mathcal{G})$. Hence the connected topological covers of the Salvetti complex are indexed by the subgroups of $\pi(\mathcal{G})$ (and so are the groupoid covers of $\mathcal{G}(\mathcal{A})$). We will use this correspondence to first, construct the covering spaces of $Sal(\mathcal{A})$ and then, to classify them.

The use of this idea can be traced back to the work of Deligne in [6], where he used this correspondence to construct the universal cover of the hyperplane complement. Then Paris generalized these arguments, using the language of oriented systems (see [18]), to show that all of the connected covers of any arrangement complement are indexed by the covers of the arrangement groupoid. Our treatment here is based on Delucchi's reformulation [7,8] of these ideas in terms of diagram of spaces. To be precise, using covers of the arrangement groupoid he defined a family of diagrams of spaces such that their homotopy colimits have the homotopy type of covers of arrangement complement. One of the benefits of his elegant treatment is that it is easy to generalize to the tangent bundle complement. This is what we plan to do now. We closely follow Delucchi's arguments as well as the notation.

Let $\mathcal{A}$ denote a submanifold arrangement in some manifold X. Let $Sal(\mathcal{A})$ be the associated Salvetti complex, the context will decide whether we view it as a CW-complex or as simplicial complex. Finally, let $\mathcal{G}$ denote the arrangement groupoid. When dealing with morphisms in $\mathcal{G}$ we will use the path-terminology. For example, domain of a morphism $[\alpha]$ will be denoted by $s(\alpha)$. Hence whenever we make statement about a path it is actually about its equivalence class in $\mathcal{G}$.

For the convenience of the reader we quickly recall relevant facts about the cover of a groupoid. Given $z \in Ob(\mathcal{G})$ its *star* is the following set of morphisms-

$$St(z) = \{[\alpha] \in Mor(\mathcal{G}) \mid s(\alpha) = z\}.$$

A *covering* of $\mathcal{G}$ is a functor $\rho\colon \mathcal{G}_\rho \to \mathcal{G}$ such that for every $z \in Ob(\mathcal{G}_\rho)$, the induced map

$$\rho_z\colon St(z) \to St(\rho(z))$$

is bijective. Given $[\alpha]$, a morphism in $\mathcal{G}$ and any $z \in \rho^{-1}(s(\alpha))$, the lift of α at z is the morphism $\rho_z^{-1}(\alpha)$, denoted by $\alpha^{<z>}$ when the covering ρ is understood. Recall that according to the main theorem of [2, Chapter 10] each object of $\mathcal{G}_\rho$ is a right coset of homotopy class of paths in the arrangement graph. Let $\mu(C \to D, F)$ denote a minimal positive path starting at a vertex $[C, C]$ ending at another vertex $[D, D]$ and it traverses an edge contained in the boundary of $[F, C]$. It is assumed that $C \prec F$.

Lemma 5.10. *With the same notation as above, $\mu(C \to D, F) \overset{+}{\sim} \mu(C \to D, F')$ if there exists $G \in \mathcal{F}^*$ such that on of the following is satisfied:*

1. *$F, F' \prec G$ and $\mu(C \to D, F), \mu(C \to D, G)$ are in the boundary of $[G, C]$;*
2. *$D = G * C$.*

Proof. Follows from Corollary 4.16. □

Definition 5.11. Given a cover of the arrangement groupoid $\rho\colon \mathcal{G}_\rho \to \mathcal{G}(\mathcal{A})$, we define a *diagram of posets* $\mathcal{D}_\rho$ on the dual face poset $\mathcal{F}^*$ as

$$\mathcal{D}_\rho(F^i) := \{v \in Ob(\mathcal{G}_\rho) \mid \rho(v) \prec F^i\}.$$

Each poset is endowed with the trivial order relation. The maps between these spaces are the following inclusions

$$\begin{array}{rcll} \mathcal{D}_\rho(F^i \to F^j)\colon \mathcal{D}(F^i) & \to & \mathcal{D}(F^j) & (i > j) \\ v & \mapsto & t(\mu(\rho(v) \to F^j \circ \rho(v), F^i)^{<v>}) & \end{array}$$

where $\mu(\rho(v) \to F^j \circ \rho(v), F^i)^{<v>}$ is the lift of the minimal positive path in $\mathcal{G}$ that starts at $[\rho(v), \rho(v)]$ and traverses an edge contained in $[F^i, \rho(v)]$.

According to the *simplicial model lemma* (Lemma 1.15), the homotopy colimit of the above diagram is in fact an order complex. We will denote this order complex by $\mathcal{S}_\rho$, its vertex set is

$$\{(F,v) \in \mathcal{F}^* \times Ob(\mathcal{G}_\rho) \mid \rho(v) \prec F\}$$

the simplices of $\mathcal{S}_\rho$ are chains with respect to the following partial order

$$(F_2, v_2) \prec_\rho (F_1, v_1) \Leftrightarrow \begin{cases} F_2 \prec F_1 \text{ and} \\ v_2 = t(\mu(\rho(v_1) \to F_2 \circ \rho(v_1), F^1)^{<v_1>}) \end{cases}$$

Remark 5.12. If ρ is the identity map then $\mathcal{S}_{id}$ is the Salvetti complex (Theorem 4.17).

Remark 5.13. Using the same arguments as in the proof of Lemma 3.11 it follows that for every chain in $\mathcal{S}_\rho$ there correspondence a chain and a vertex in $\mathcal{F}^*$. If ϕ is a chain in $\mathcal{F}^*$ and $v \in Ob(\mathcal{G}_\rho)$ such that $\rho(v) \prec max(\phi)$, then the pair (ϕ, v) corresponds to a chain in $\mathcal{S}_\rho$ and every chain is of this form. Moreover, the simplicial complex $\Delta(\mathcal{S}_\rho)$ is the barycentric subdivision of a CW-complex $\mathcal{S}_\rho^{CW}$; the barycentric subdivision of a (closed) k-cell of this CW-complex, which we denote by a pair $[F^k, v]$, is given by

$$[F^k, v] := \bigcup_{max(\phi)=F^k} (\phi, v).$$

Vertices of $\mathcal{S}_\rho^{CW}$ are of the form $[\rho(v), v]$, hence are in bijection with the objects of $\mathcal{G}_\rho$. For a k-cell $[F^k, v]$ a 0-cell $[\rho(v'), v']$ is its vertex if and only if

$$v' = t(\mu(\rho(v) \to F' \circ \rho(v'), F^k)^{<v>})$$

for some $F' \prec F^k$.

Our aim is to show that the simplicial complex $\mathcal{S}_\rho$ is a covering space of the (simplicial) Salvetti complex. The strategy we employ here is motivated by the proof of a well known fact in elementary algebraic topology. A simplicial and fixed point free action of a discrete group on a simplicial complex is a covering space action (see for example [13, Section 1.B]). First, we prove that there is a natural, simplicial map between these two complexes and then proceed on to prove that it is indeed a covering map.

The functor $\rho\colon \mathcal{G}_\rho \to \mathcal{G}$ naturally induces a morphism of diagrams $\lambda\colon \mathcal{D}_\rho \to \mathcal{D}_{id}$. On the level of spaces it is given by:

$$\begin{array}{rcll} \lambda_F\colon \mathcal{D}_\rho(F) & \to & \mathcal{D}_{id}(F) & \forall F \in \mathcal{F}^* \\ v & \mapsto & \rho(v). & \end{array}$$

In order to check the compatibility of this morphism with the maps between the spaces, consider $F_2 \prec F_1$ and $v \in \mathcal{D}_\rho(F_2)$. Then,

$$\begin{array}{rcl} \lambda_{F_2}(\mathcal{D}_\rho(F_1 \to F_2)(v)) & = & \rho(t(\mu(\rho(v) \to F_2 \circ \rho(v), F_1)^{<v>})) \\ & = & F_2 \circ \rho(v) \\ & = & \mathcal{D}_{id}(F_1 \to F_2)(\rho(v)) \\ & = & \mathcal{D}_{id}(F_1 \to F_2)(\lambda_{F_1}(v)). \end{array}$$

The morphism λ induces a map between homotopy colimits by functoriality, hence a map of simplicial complexes

$$\Lambda_\rho\colon \mathcal{S}_\rho \to Sal(\mathcal{A}).$$

Λ_ρ maps the simplex (ϕ, v) of $\mathcal{S}_\rho$ to the simplex $(\phi, \rho(v))$ of $Sal(\mathcal{A})$. Moreover, a morphism $\eta\colon \mathcal{G}_{\rho_1} \to \mathcal{G}_{\rho_2}$ between two covers of the arrangement groupoids induces a map $\Lambda_\eta\colon \mathcal{S}_{\rho_1} \to \mathcal{S}_{\rho_2}$.

Theorem 5.14. *The map $\Lambda_\rho\colon \mathcal{S}_\rho \to Sal(\mathcal{A})$, induced by a cover $\rho\colon \mathcal{G}_\rho \to \mathcal{G}$ of the arrangement groupoid, is a covering map.*

Proof. First note that $Sal(\mathcal{A})$ is a finite dimensional and locally finite simplicial complex implying that it is connected and locally path connected. Hence all its connected covering spaces exist.

Let $p \in Sal(\mathcal{A})$ be some point, σ be the smallest dimensional simplex containing p, let U denote the star of σ (set of all simplices of $Sal(\mathcal{A})$ that contain σ). In order to show that Λ_ρ is a covering map it is enough to show that $\Lambda_\rho^{-1}(U)$ is an even covering of U.

Let us first characterize U and its inverse image in terms of chains and chambers. As σ is a simplex of $Sal(\mathcal{A})$ there exists a chain $\tilde{\phi}$ in $\mathcal{F}^*$ and (a chamber) $\tilde{C} \prec max(\tilde{\phi}) =: \tilde{F}$ such that $\sigma = (\tilde{\phi}, \tilde{C})$. Thus star of σ takes the following form:

$$U = \bigcup_{\substack{\phi \supseteq \tilde{\phi} \\ C \in R(\phi, \tilde{C})}} (\phi, C)$$

where $R(\phi, \tilde{C}) := \{C \prec max(\phi) \mid \tilde{F} \circ C = \tilde{C}\}$. As a set, the pre-image of U is:

$$\Lambda_\rho^{-1}(U) = \bigcup_{\substack{\phi \supseteq \tilde{\phi} \\ \{v \ \mid \ \rho(v) \in R(\phi, \tilde{C})\}}} (\phi, v).$$

For a fixed object $w \in \rho^{-1}(\tilde{C})$ define

$$W_w := \bigcup_{\substack{\phi \supseteq \tilde{\phi} \\ C \in R(\phi, \tilde{C})}} (\phi, v(C, w))$$

where $v(C, w)$ is the (unique) object, of $\mathcal{G}_\rho$, in $\rho^{-1}(C)$ from which w can be reached by the lift of a minimal positive path.

Rest of the proof follows from the following three claims which are left as an exercise.

Claim 1: For any $w_1 \neq w_2 \in \rho^{-1}(\tilde{C})$, $W_{w_1} \cap W_{w_2} = \emptyset$.

Claim 2:

$$\coprod_{w \in \rho^{-1}(\tilde{C})} W_w = \Lambda_\rho^{-1}(U).$$

Claim 3: Fix $w \in \rho^{-1}(\tilde{C})$, then $\Lambda_\rho\colon W_w \to U$ is a homeomorphism. □

Example 5.15. Consider the arrangement $\mathcal{A}_1$ of 2 points in S^1 (Example 2.10), its associated Salvetti complex $Sal(\mathcal{A}_1)$ was described in Figure 6. Figure 9 depicts a portion of its universal cover. The objects of the covering groupoid are denoted by $\tilde{A}, \tilde{B}$ (indices are suppressed for notational simplicity). Note that there are two minimal positive paths, in $Sal(\mathcal{A})$, from $[B, B]$ to $[A, A]$. One of them is the (oriented) edge $[p, B]$ and the other is the edge $[q, B]$. Both of these paths are represent distinct equivalence classes in $\mathcal{G}^+(A)$ as well as $\mathcal{G}(A)$. In the covering space, end points of the lifts of these two paths are distinct and are contained in $\rho^{-1}([A, A])$. Also note that the functor $J\colon \mathcal{G}^+(A) \to \mathcal{G}(A)$ is faithful in this example.

Now we prove that any connected topological covering space of the Salvetti complex corresponds to a Salvetti-type diagram model. We use the correspondence between isomorphism classes of covering spaces and conjugacy classes of subgroups ([13, Theorem 1.38]).



FIGURE 9. The Universal cover of $Sal(\mathcal{A}_1)$

Given a cover $\rho\colon \mathcal{G}_\rho \to \mathcal{G}$ of the arrangement groupoid $\mathcal{G}$, consider the corresponding covering cell complex $\mathcal{S}_\rho$ of the associated Salvetti complex $Sal(\mathcal{A})$ (see Remark 5.13).
2-skeleton: For a codimension 2 face F^2 and an object $v \in Ob(\mathcal{G}_\rho)$ (such that $\rho(v) \prec F^2$) there is one 2-cell $[F^2, v]$ in $\mathcal{S}_\rho$. The vertices of $[F^2, v]$ are indexed by the chambers whose closures contain F^2. To such a chamber C, there corresponds an object in the covering groupoid given by $t(\rho(v) \to C, F^2)^{<v>}$. Hence the vertex set of $[F^2, v]$ is given by $\{[\rho(v_i), v_i] \mid \rho(v_i) \prec F^2, i = 1, \ldots, 2k\}$. A 1-cell $[F^1, v_i]$ in the boundary of $[F^2, v]$ corresponds to codimension 1 face F^1 such that $F^1 \prec F^2$. The vertex v_1 is such that the chamber $\rho(v_i)$ is on the same side as $\rho(v)$ with respect to F^1. Moreover, the cell is oriented to have v_i as its start vertex.

Remark 5.16. Attachment of higher dimensional cells is analogous to the construction of Salvetti complex. Hence it can be shown that these covering complexes $\mathcal{S}_\rho$ have the structure of a MH-complex.

The next step in the classification is to show that the fundamental group of $\mathcal{S}_\rho$ is isomorphic to the object group of $\mathcal{G}_\rho$. We compare the relations given by 2-cells of $\mathcal{S}_\rho$ and the relations that define $\mathcal{G}_\rho$.

Recall that the equivalence relation in the arrangement groupoid $\mathcal{G}$ is generated by identifying two paths with same end points which form the boundary of a 2-cell in $Sal(\mathcal{A})$. As a result, the identity in an object group is an equivalence class of loops based at that object that can be written as $\gamma\gamma'^{-1}$, where γ, γ' are paths bounding a 2-cell. For a given groupoid cover $\rho\colon \mathcal{G}_\rho \to \mathcal{G}$ let G_ρ denote the graph underlying $\mathcal{G}_\rho$. The morphisms of $\mathcal{G}_\rho$ are equivalence classes of paths in the graph G_ρ. This equivalence relation is the one induced by ρ. To be precise, two paths $\tilde{\alpha}, \tilde{\beta}$ in G_ρ are identified if and only if they have same end points and $\rho(\tilde{\alpha}) \sim \rho(\tilde{\beta})$ (where $\tilde{\alpha}, \tilde{\beta}$ are lifts of α, β respectively). Hence the relations in $\mathcal{G}_\rho$ are generated by the loops of the form $\tilde{\gamma_1}\tilde{\gamma_2}^{-1}$ such that $[\gamma_1] = [\gamma_2]$ (in $\mathcal{G}_\rho$), denote the normal subgroup generated by these loops by Σ_ρ. In a nutshell we have $\pi(\mathcal{G}_\rho) \cong \pi_1(G_\rho)/\Sigma_\rho$.

On the other hand let σ_ρ denote the smallest normal subgroup of $\pi_1((\mathcal{S}_\rho)_1)$ generated by the relations imposed by the boundary of every 2-cell. Then, $\pi_1(\mathcal{S}_\rho) \cong \pi_1((\mathcal{S}_\rho)_1)/\sigma_\rho$.

Theorem 5.17. *With the notation as above:*

$$\pi(\mathcal{G}_\rho) \cong \pi_1(\mathcal{S}_\rho).$$

Now we are in the position to state the classification of covering spaces of the Salvetti complex.

Theorem 5.18. *Let $\mathcal{A}$ be a submanifold arrangement in a l-manifold X, let $Sal(\mathcal{A})$ denote the associated Salvetti complex and $\mathcal{G}$ denote the arrangement groupoid. For any topological cover $p\colon S \to Sal(\mathcal{A})$, there exists a cover of the arrangement groupoid $\rho\colon \mathcal{G}_\rho \to \mathcal{G}$ such that the homotopy colimit of the associated diagram of spaces $\mathcal{D}_\rho$ is isomorphic to S as a covering space of $Sal(\mathcal{A})$*

Proof. Let H denote the isomorphic image of $p_*(\pi_1(S))$ inside $\pi(\mathcal{G}(\mathcal{A}))$ (under the isomorphism in Theorem 5.17). As a result of [2, Theorem 10.6.1] we see that there is a covering groupoid $\rho\colon \mathcal{G}_\rho \to \mathcal{G}(A)$ such that $\pi(\mathcal{G}_\rho) \cong H$. Let S_ρ denote the homotopy colimit of the diagram of spaces defined using $\mathcal{G}_\rho$. Again appealing to Theorem 5.17 it implies that $\pi_1(S_\rho) \cong H$. Let ι_ρ denote the inclusion of G_ρ (the graph underlying $\mathcal{G}_\rho$) into S_ρ as its 1-skeleton. Since $\mathcal{G}_\rho$ is equivalent to the fundamental groupoid of S_ρ the following diagram commutes.

$$\begin{tikzcd}
\pi(\mathcal{G}_\rho) \arrow[r, "\cong"] \arrow[d, "\rho_*"'] & \pi_1(S_\rho) \arrow[d, "(\Lambda_\rho)_*"] \\
\pi(\mathcal{G}) \arrow[r, "\cong"] & \pi_1(Sal(\mathcal{A}))
\end{tikzcd}$$

Since $(\Lambda_\rho)_*(S_\rho) \cong H$ we have that S and S_ρ are isomorphic as covering spaces. $\square$

Corollary 5.19. *Any cover $p\colon S \to Sal(\mathcal{A})$ of the Salvetti complex can be realized as the order complex of a poset.*

Corollary 5.20. *Let $\hat{\rho}\colon \mathcal{G}_{\hat{\rho}} \to \mathcal{G}$ denote the universal cover of the arrangement groupoid. Then $\mathcal{S}_{\hat{\rho}}$, the homotopy colimit of the associated diagram of spaces (posets) is the universal cover of $Sal(\mathcal{A})$.*

Concluding Remarks

One of the important topological invariants not mentioned in this paper is the cohomology ring of the tangent bundle complement. For hyperplane arrangements there is a beautiful theory of cohomology of the complement (known as the Orlik-Solomon algebra, see [17] for details). However for submanifold arrangements there is very little understanding of how the combinatorics determines the cohomology ring. It is a work in progress. However, we would like to point the reader to [9, Section 3.7] for some calculations of cohomology groups using the Bousfield-Kan spectral sequence. It is conjectured that there is a finer grading of cohomology groups indexed by the intersection data (see [9, Conjecture 3.7.8]).

As an application of the theory developed so far it is possible study toric arrangements. Recently there has been an intense activity in this field we refer the reader to [9, Section 4.2] for a survey of this subject and some open questions.

Deligne's work [6] is still influential in the topological study of finite Coxeter groups and associated Artin groups. Similar to spherical reflection groups there is a theory of geometric reflection groups and more generally manifold reflection groups, for example, see [4, Chapters 6, 10] and [12]. The fixed points of the action of such reflection group on a manifold is a submanifold arrangement. The fundamental group of the associated tangent bundle complement can be thought of as a generalization of Artin groups. It is a work in progress to understand these groups using submanifold arrangements.

REFERENCES

[1] A. Björner and G. M. Ziegler. Combinatorial stratification of complex arrangements. *J. Amer. Math. Soc.*, 5(1):105–149, 1992. 2, 22

[2] R. Brown. *Topology and groupoids*. BookSurge, LLC, Charleston, SC, 2006. 26, 27, 31

[3] R. Charney. Artin groups of finite type are biautomatic. *Math. Ann.*, 292(4):671–683, 1992. 2

[4] M. W. Davis. *The geometry and topology of Coxeter groups*, volume 32 of *London Mathematical Society Monographs Series*. Princeton University Press, Princeton, NJ, 2008. 31

[5] P. Dehornoy and L. Paris. Gaussian groups and Garside groups, two generalisations of Artin groups. *Proc. London Math. Soc. (3)*, 79(3):569–604, 1999. 2

[6] P. Deligne. Les immeubles des groupes de tresses généralisés. *Invent. Math.*, 17:273–302, 1972. 2, 16, 24, 26, 31

[7] E. Delucchi. *Topology and combinatorics of arrangement covers and of nested set complexes*. PhD thesis, ETH Zürich, 2006. 1, 2, 24, 27

[8] E. Delucchi. Combinatorics of covers of complexified hyperplane arrangements. *Progress in Mathematics*, 283:138, 2007. 27

[9] P. Deshpande. *Arrangements of Submanifolds and the Tangent Bundle Complement*. PhD thesis, The University of Western Ontario, 2011. Electronic Thesis and Dissertation Repository. `http://ir.lib.uwo.ca/etd/154`. 3, 8, 9, 26, 31

[10] P. Gabriel and M. Zisman. *Calculus of fractions and homotopy theory*. Springer-Verlag, 1967. 25

[11] F. A. Garside. The braid group and other groups. *Quart. J. Math. Oxford Ser. (2)*, 20:235–254, 1969. 2

[12] E. Gutkin. Geometry and combinatorics of groups generated by reflections. *Enseign. Math. (2)*, 32(1-2):95–110, 1986. 31

[13] A. Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002. 28, 29

[14] P. J. Higgins. *Notes on categories and groupoids*. Van Nostrand Reinhold Co., London, 1971. Van Nostrand Rienhold Mathematical Studies, No. 32. 24, 26

[15] D. Kozlov. *Combinatorial algebraic topology*, volume 21 of *Algorithms and Computation in Mathematics*. Springer, Berlin, 2008. 4, 5

[16] L. Moci. Combinatorics and topology of toric arrangements defined by root systems. *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.*, 19(4):293–308, 2008. 9

[17] P. Orlik and H. Terao. *Arrangements of hyperplanes*, volume 300 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1992. 4, 31

[18] L. Paris. The covers of a complexified real arrangement of hyperplanes and their fundamental groups. *Topology Appl.*, 53(1):75–103, 1993. 24, 26

[19] L. Paris. Universal cover of salvetti's complex and topology of simplicial arrangements of hyperplanes. *Transactions of the American Mathematical Society*, 340(1):149–178, Nov. 1993. 1, 2, 4, 14

[20] T. B. Rushing. *Topological embeddings*. Academic Press, New York, 1973. Pure and Applied Mathematics, Vol. 52. 7, 8, 10

[21] M. Salvetti. Topology of the complement of real hyperplanes in N. *Inventiones Mathematicae*, 88(3):603–618, Oct. 1987. 1, 2, 4, 17, 18

[22] M. Salvetti. On the homotopy theory of complexes associated to metrical-hemisphere complexes. *Discrete Math.*, 113(1-3):155–177, 1993. 2, 20, 22, 26

[23] R. Scott. Blow-ups and complements in zonotopal complexes, Oct. 2004.
Available at `http://www.msri.org/web/msri/online-videos/-/video/showVideo/1837`. 2

[24] V. Welker, G. M. Ziegler, and R. T. Živaljević. Homotopy colimits - comparison lemmas for combinatorial applications. *Journal für die Reine und Angewandte Mathematik*, 509:117–149, 1999. 5, 6, 7

[25] G. M. Ziegler. *Lectures on polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. 19

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MA
E-mail address: `p.deshpande@neu.edu`